

Archives of Biochemical Engineering

Review Article

Comparative Analysis of enzyme immobilization techniques versus Standard Care in low bioreactor yield

Yoshida N^{1*}, Suzuki K² and Ito S³

¹Department of Bioprocess Technology, Ganges Academy of Biochemical Engineering, Malaysia

²Department of Bioprocess Technology, Euphrates Institute of Bioprocess Technology, Japan

³Department of Biochemical Engineering, Euphrates Institute of Bioprocess Technology, Taiwan

*Address for Correspondence: Naomi Yoshida, Department of Bioprocess Technology, Ganges Academy of Biochemical Engineering, Malaysia; E-mail: naomi.yoshida@ganges.edu

Received: 15 October 2025; Accepted: 5 January 2026; Published: 2 April 2026.

Citation of this article: Yoshida N, Suzuki K, Ito S (2026) Comparative Analysis of enzyme immobilization techniques versus Standard Care in low bioreactor yield. Arch Biochem Eng, 1(1): 01-06. DOI: [10.5281/zenodo.22812454](https://doi.org/10.5281/zenodo.22812454)

Copyright: © 2026 Naomi Yoshida, et al. This is an open access article distributed under the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Low volumetric yield in bioreactors run with free, dissolved biocatalyst -- the standard-care configuration in most industrial enzymatic processes -- is driven by enzyme deactivation, product inhibition, and the cost of discarding a soluble catalyst after a single batch. This comparative analysis reviews four widely used immobilization strategies -- adsorption, covalent binding, entrapment/encapsulation, and cross-linked enzyme aggregates (CLEAs) -- against free-enzyme standard care, focused on reaction yield, operational stability across reuse cycles, and volumetric productivity under diffusion-limited bioreactor conditions.

Immobilized preparations consistently retain a larger fraction of initial activity after repeated batches than free enzyme, and packed-bed or fluidized-bed configurations built around immobilized catalyst achieve markedly higher volumetric productivity than standard-care stirred-tank operation limited by single-use enzyme. CLEAs and covalent binding show the strongest combination of activity retention and yield recovery, while adsorption and entrapment trade some activity for simpler, lower-cost preparation; diffusional resistance from the support remains the dominant yield-limiting factor and must be weighed against the reusability gains immobilization provides. Immobilization is therefore not a uniform upgrade over standard care but a family of trade-offs whose optimal choice depends on the specific low-yield bottleneck, with reactor configuration selected jointly with immobilization technique rather than as an independent decision.

Keywords: enzyme immobilization; bioreactor yield; cross-linked enzyme aggregates; operational stability; biocatalysis

1. Introduction

Free, dissolved enzyme run once through a stirred-tank batch and then discarded remains the default "standard care" configuration in much of industrial biocatalysis, largely because it needs no carrier development before a process can start. It is also the configuration most exposed to the causes of low bioreactor yield: progressive catalyst deactivation, product inhibition that a single-pass reactor cannot mitigate, and the irrecoverable loss of active enzyme at the end of every batch, which raises effective catalyst cost per unit product to a level that can make an otherwise viable bioprocess uneconomic [1,2].

Enzyme immobilization -- fixing the biocatalyst to or within an insoluble support so it can be recovered and reused -- has been proposed as a remedy since at least the 1970s: an immobilized catalyst that survives ten reuse cycles at 70% of initial activity delivers far more product per unit of enzyme purchased than a free catalyst discarded after one cycle, even at a somewhat lower per-cycle yield [3,4]. Adsorption, covalent binding, entrapment/encapsulation, and cross-linked enzyme aggregates (CLEAs) each trade preparation simplicity, activity retention, and diffusional resistance differently, and a process engineer choosing among them, or against standard care, needs those trade-offs stated against the yield metrics that determine bioreactor economics [5,6].

This analysis compares four immobilization strategies against free-enzyme standard care across the three determinants of realized bioreactor yield that recur in the literature: intrinsic reaction yield, operational stability across repeated batches, and the diffusional penalty a support imposes at bioreactor scale. The aim is to state which low-yield bottleneck each technique addresses, since no single method dominates standard care, or every other method, across all contexts [7,8].

2. Materials and Methods

This analysis synthesizes the major review syntheses on enzyme immobilization technique and industrial

application, supplemented by primary comparative and mechanistic studies on the techniques reviewed [9,10,11,12,13,14]. Four techniques were selected because they span the principal mechanisms by which a biocatalyst is rendered insoluble and reusable: physical adsorption, covalent attachment, entrapment/encapsulation, and carrier-free cross-linking into CLEAs [15,16,17,18,19].

For each technique, three categories of evidence were extracted where reported: relative activity and yield immediately after immobilization against the free-enzyme baseline; residual activity after repeated reuse cycles; and volumetric productivity under bioreactor-relevant, diffusion-influenced conditions rather than idealized assay conditions [7,20,21]. CLEA evidence was weighted toward the founding methodology papers and subsequent cross-linker optimization studies, since cross-linking density is a major determinant of both activity retention and diffusional resistance [22,23,24,25]. Process-economic evidence on preparation cost, reusability, and scale-up was drawn from the biocatalytic process-engineering literature on when immobilization becomes cost-justified relative to single-use free enzyme [26,27,28,29], and mass-transfer considerations general to bioreactor scale-up were drawn from the bioreactor engineering literature [30,31].

Figures and tables below summarize representative, illustrative values drawn from the ranges reported across these sources; where a range rather than a point estimate is reported, a representative midpoint is used, and the rank order of technique performance, not any single numeric value, is what the underlying literature supports most consistently.

3. Comparative catalytic performance and reaction yield

Every technique reviewed loses some intrinsic activity relative to free enzyme in solution immediately after immobilization, because attachment restricts

conformational flexibility, can distort the active site, and introduces a diffusional barrier even before any reuse-related deactivation [9,4]. Adsorption, relying on weak reversible interactions, typically preserves the largest fraction of native activity, but that same weak binding permits leaching and an erosion of apparent yield over successive batches that is easily mistaken for catalytic deactivation [13,12].

Table 1: Representative initial reaction yield and principal trade-off for free enzyme (standard care) and four immobilization techniques, drawn from comparative ranges reported across the reviewed literature [9,13,19].			
Preparation	Reaction yield vs. free enzyme	Leaching risk	Principal trade-off
Free enzyme (standard care)	100% (reference)	N/A - not reusable	No reuse; catalyst discarded each batch
Adsorption	70-90%	High	Simple, low-cost; activity lost to leaching over time
Covalent binding	60-85%	Very low	Stable attachment; coupling can hinder active site
Entrapment/encapsulation	50-75%	Low	Preserves native structure; strong diffusional resistance
Cross-linked enzyme aggregates	70-95%	Very low	No carrier needed; cross-linking density must be optimized

Covalent immobilization essentially eliminates leaching, but coupling through lysine, cysteine, or other reactive side chains can reduce initial activity more than adsorption if binding occurs near the active site, a problem multi-point attachment protocols address by favoring surface regions distant from the catalytic center [9,32]. Entrapment/encapsulation preserves native activity at the molecular level but suffers the largest

apparent yield penalty in practice, since substrate and product must diffuse through the matrix [18,17]. Naturally derived biopolymer carriers such as chitosan and alginate partially offset this penalty through low cost at production scale [33], and carrier-free architectures such as protein-inorganic hybrid nanoflowers have in some assays exceeded free-enzyme activity, showing the trade-off is not absolute across every emerging technique [34].

Table 1 summarizes representative reaction-yield outcomes against free-enzyme standard care. CLEAs, requiring no exogenous carrier, avoid both the carrier-dilution problem of carrier-bound preparations and much of adsorption's leaching problem, with activity retention that in several studies approaches or exceeds covalent preparations at lower preparation cost [19,22,23].

4. Operational stability and reusability

The largest source of yield loss under standard care is not per-batch inefficiency but that the catalyst cannot be reused: every batch begins with fresh enzyme and ends with it discarded, so realized yield per unit enzyme purchased is bounded by a single cycle regardless of how well that cycle performs [1,2]. Operational-stability data -- residual activity across reuse cycles -- is where immobilized preparations show their most consistent advantage over standard care [8,7].

Figure 1 summarizes representative residual activity after ten reuse cycles. Free enzyme, not reusable by construction, retains almost nothing; the immobilized preparations retain a substantial, technique-dependent fraction. Covalent binding and CLEAs show the strongest reuse performance, consistent with their resistance to leaching and, for CLEAs, the rigidity conferred by cross-linking, while adsorption shows more variable retention that depends on carrier chemistry and process ionic strength [25,24]. Entrapped preparations retain intrinsic activity well but at the cost of a lower activity ceiling from diffusional resistance [18].




Figure 1: Representative residual catalytic activity after ten reuse cycles for free enzyme (standard care) and four immobilization techniques, illustrating the reuse-stability advantage of covalent binding and cross-linked enzyme aggregates over adsorption, entrapment and standard care [8,25,19].

5. Diffusional limitations and bioreactor-scale yield

Reuse stability alone does not guarantee superior bioreactor-scale yield, because immobilization introduces a mass-transfer penalty that dissolved standard-care enzyme does not face. Substrate and product must diffuse to and from the immobilized site, competing kinetically with catalytic turnover; when diffusion is slow relative to turnover the reaction becomes diffusion-limited, a problem worse for larger particles, denser cross-linking, and more viscous media [4,21].

This is the tension that determines which configuration resolves a genuinely low-yield process: standard care avoids diffusional resistance but pays with catalyst loss, while immobilized preparations recover catalyst but must be engineered -- particle size, porosity, reactor configuration -- to keep diffusion from eroding that gain [15,20]. Figure 2 summarizes representative volumetric productivity across configurations. Packed-bed and fluidized-bed reactors built around immobilized catalyst consistently outperform standard-care stirred-tank operation, because continuous operation with a retained catalyst bed sustains high effective loading over time that a single-use batch cannot match, even though the per-particle reaction is itself diffusion-limited [28,27]. Mass-transfer coefficients generally worsen on scale-up, which penalizes any configuration, standard care included, lacking a mechanism to sustain high local

catalyst concentration near substrate [30,31].




Figure 2: Representative volumetric productivity by bioreactor configuration, comparing free-enzyme standard-care stirred-tank operation with packed-bed, fluidized-bed and retained-catalyst stirred-tank configurations built around immobilized enzyme [28,20,30].

6. Discussion

Table 2 draws together reusability, relative cost, and scale-up character to frame the practical choice a process engineer faces. The evidence does not support treating immobilization as a uniform upgrade over standard care; rather it supports matching technique to the specific bottleneck responsible for the low yield observed [7,14]. Where catalyst instability and single-use cost dominate, CLEAs and covalent binding offer the strongest combination of retention and leaching resistance, and their reuse economics justify the added complexity once more than a handful of cycles are achievable [19,26]. Where the reaction tolerates a moderate diffusional penalty and rapid, low-cost deployment matters more than maximal reuse, adsorption or entrapment may match or exceed the economics of covalent immobilization and standard care without cross-linking or coupling overhead [13,17].

Technique and reactor configuration are not independent choices. A CLEA or covalently bound catalyst with excellent reuse stability delivers little bioreactor-scale benefit if still run as a stirred-tank batch that discards catalyst between runs; the yield gains reported for immobilized systems are realized specifically when immobilization is paired with a configuration --

packed-bed, fluidized-bed, or retained-catalyst stirred tank -- engineered to exploit catalyst recovery [28,20]. Conversely, a packed-bed or fluidized-bed configuration built on a poorly stable adsorbed enzyme can foul or lose activity faster than its improved mass-transfer geometry compensates for [25,6]. The economic case for moving away from standard care rests on the joint choice of technique and configuration evaluated against yield bottleneck, enzyme cost, and achievable reuse cycles, not on any single technique's laboratory-scale figures taken in isolation [27,29].

Table 2: Reusability, relative preparation cost and scale-up character of free enzyme (standard care) and four immobilization techniques [26,27,28].

Preparation	Typical reuse cycles	Relative preparation cost	Preferred reactor configuration
Free enzyme (standard care)	1 (single use)	Lowest	Stirred-tank batch, single pass
Adsorption	Low-moderate	Low	Packed-bed or stirred-tank, short campaigns
Covalent binding	High	Moderate-high	Packed-bed or fluidized-bed
Entrapment/enapsulation	Moderate-high	Moderate	Fluidized-bed or membrane reactor
Cross-linked enzyme aggregates	High	Moderate	Packed-bed or retained-catalyst stirred-tank

7. Conclusion

Standard care remains the simplest configuration to deploy but is structurally exposed to the two dominant causes of low bioreactor yield reviewed here: the impossibility of catalyst reuse and the resulting cap on productivity per unit enzyme purchased. Immobilization addresses catalyst reuse and, for several techniques, substantially improves both operational stability and bioreactor-scale productivity relative to standard care, but every technique trades some initial activity for a diffusional resistance that must be engineered around [1,4]. CLEAs and covalent binding offer the strongest overall combination of activity retention, leaching resistance, and reuse-driven yield recovery; adsorption

and entrapment remain competitive where preparation simplicity or diffusional tolerance outweighs maximal reuse; and in every case the realized yield benefit depends on pairing technique with a reactor configuration engineered to exploit catalyst recovery [19,28,7]. The choice between standard care and a specific immobilization route, and the reactor configuration paired with it, should be treated as a single joint decision driven by the process's instability, mass-transfer, and reuse-economics profile.

8. References

- Sheldon, R.A. (2007) Enzyme immobilization: the quest for optimum performance. *Advanced Synthesis & Catalysis*, 349(8-9): 1289-1307. DOI: [10.1002/adsc.200700082](https://doi.org/10.1002/adsc.200700082)
- Woodley, J.M. (2008) New opportunities for biocatalysis: making pharmaceutical processes greener. *Trends in Biotechnology*, 26(6): 321-327. DOI: [10.1016/j.tibtech.2008.03.004](https://doi.org/10.1016/j.tibtech.2008.03.004)
- Klibanov, A.M. (1983) Immobilized enzymes and cells as practical catalysts. *Science*, 219(4585): 722-727. DOI: [10.1126/science.219.4585.722](https://doi.org/10.1126/science.219.4585.722)
- Cao, L., van Langen, L., Sheldon, R.A. (2003) Immobilised enzymes: carrier-bound or carrier-free? *Current Opinion in Biotechnology*, 14(4): 387-394. DOI: [10.1016/S0958-1669\(03\)00096-X](https://doi.org/10.1016/S0958-1669(03)00096-X)
- Sheldon, R.A., van Pelt, S. (2013) Enzyme immobilisation in biocatalysis: why, what and how. *Chemical Society Reviews*, 42(15): 6223-6235. DOI: [10.1039/c3cs60075k](https://doi.org/10.1039/c3cs60075k)
- Hanefeld, U., Gardossi, L., Magner, E. (2009) Understanding enzyme immobilisation. *Chemical Society Reviews*, 38(2): 453-468. DOI: [10.1039/b711564b](https://doi.org/10.1039/b711564b)
- Garcia-Galan, C., Berenguer-Murcia, A., Fernandez-Lafuente, R., Rodrigues, R.C. (2011) Potential of different enzyme immobilization strategies to improve enzyme performance. *Advanced Synthesis & Catalysis*, 353(16): 2885-2904. DOI: [10.1002/adsc.201100288](https://doi.org/10.1002/adsc.201100288)

[10.1002/adsc.201100534](https://doi.org/10.1002/adsc.201100534)

8. Rodrigues, R.C., Ortiz, C., Berenguer-Murcia, A., Torres, R., Fernandez-Lafuente, R. (2013) Modifying enzyme activity and selectivity by immobilization. Chemical Society Reviews, 42(15): 6290-6307. DOI: [10.1039/c2cs35231a](https://doi.org/10.1039/c2cs35231a)
9. Mateo, C., Palomo, J.M., Fernandez-Lorente, G., Guisan, J.M., Fernandez-Lafuente, R. (2007) Improvement of enzyme activity, stability and selectivity via immobilization techniques. Enzyme and Microbial Technology, 40(6): 1451-1463. DOI: [10.1016/j.enzmictec.2007.01.018](https://doi.org/10.1016/j.enzmictec.2007.01.018)
10. Datta, S., Christena, L.R., Rajaram, Y.R.S. (2013) Enzyme immobilization: an overview on techniques and support materials. 3 Biotech, 3(1): 1-9. DOI: [10.1007/s13205-012-0071-7](https://doi.org/10.1007/s13205-012-0071-7)
11. Brady, D., Jordaan, J. (2009) Advances in enzyme immobilisation. Biotechnology Letters, 31(11): 1639-1650. DOI: [10.1007/s10529-009-0076-4](https://doi.org/10.1007/s10529-009-0076-4)
12. Homaei, A.A., Sariri, R., Vianello, F., Stevanato, R. (2013) Enzyme immobilization: an update. Journal of Chemical Biology, 6(4): 185-205. DOI: [10.1007/s12154-013-0102-9](https://doi.org/10.1007/s12154-013-0102-9)
13. Mohamad, N.R., Marzuki, N.H.C., Buang, N.A., Huyop, F., Wahab, R.A. (2015) An overview of technologies for immobilization of enzymes and surface analysis techniques for immobilized enzymes. Biotechnology & Biotechnological Equipment, 29(2): 205-220. DOI: [10.1080/13102818.2015.1008192](https://doi.org/10.1080/13102818.2015.1008192)
14. Sirisha, V.L., Jain, A., Jain, A. (2016) Enzyme immobilization: an overview on methods, support material, and applications of immobilized enzymes. Advances in Food and Nutrition Research, 79: 179-211. DOI: [10.1016/bs.afnr.2016.07.004](https://doi.org/10.1016/bs.afnr.2016.07.004)
15. Zdarta, J., Meyer, A.S., Jesionowski, T., Pinelo, M. (2018) A general overview of support materials for enzyme immobilization: characteristics, properties, practical application. Catalysts, 8(2): 92. DOI: [10.3390/catal8020092](https://doi.org/10.3390/catal8020092)
16. Jesionowski, T., Zdarta, J., Krajewska, B. (2014) Enzyme immobilization by adsorption: a review. Adsorption, 20(5-6): 801-821. DOI: [10.1007/s10450-014-9623-y](https://doi.org/10.1007/s10450-014-9623-y)
17. Krajewska, B. (2004) Application of chitin- and chitosan-based materials for enzyme immobilizations: a review. Enzyme and Microbial Technology, 35(2-3): 126-139. DOI: [10.1016/j.enzmictec.2003.12.013](https://doi.org/10.1016/j.enzmictec.2003.12.013)
18. Betancor, L., Luckarift, H.R. (2008) Bioinspired enzyme encapsulation for biocatalysis. Trends in Biotechnology, 26(10): 566-572. DOI: [10.1016/j.tibtech.2008.06.009](https://doi.org/10.1016/j.tibtech.2008.06.009)
19. Sheldon, R.A. (2011) Cross-linked enzyme aggregates as industrial biocatalysts. Organic Process Research & Development, 15(1): 213-223. DOI: [10.1021/op100289f](https://doi.org/10.1021/op100289f)
20. Cantone, S., Ferrario, V., Corici, L., Ebert, C., Fattor, D., Spizzo, P., Gardossi, L. (2013) Efficient immobilisation of industrial biocatalysts: criteria and constraints for the selection of organic polymeric carriers and immobilisation methods. Chemical Society Reviews, 42(15): 6262-6276. DOI: [10.1039/c3cs35464d](https://doi.org/10.1039/c3cs35464d)
21. Liese, A., Hilterhaus, L. (2013) Evaluation of immobilized enzymes for industrial applications. Chemical Society Reviews, 42(15): 6236-6249. DOI: [10.1039/c3cs35511j](https://doi.org/10.1039/c3cs35511j)
22. Cao, L., van Rantwijk, F., Sheldon, R.A. (2000) Cross-linked enzyme aggregates: a simple and effective method for the immobilization of penicillin acylase. Organic Letters, 2(10): 1361-1364. DOI: [10.1021/ol005593x](https://doi.org/10.1021/ol005593x)
23. Schoevaart, R., Wolbers, M.W., Golubovic, M., Ottens, M., Kieboom, A.P.G., van Rantwijk, F., van der Wielen, L.A.M., Sheldon, R.A. (2004) Preparation, optimization, and structures of cross-linked enzyme aggregates (CLEAs). Biotechnology and Bioengineering, 87(6): 754-762. DOI: [10.1002/bit.20184](https://doi.org/10.1002/bit.20184)
24. Barbosa, O., Ortiz, C., Berenguer-Murcia, A., Torres, R., Rodrigues, R.C., Fernandez-Lafuente, R. (2014) Glutaraldehyde in bio-catalysts design: a useful crosslinker and a versatile tool in enzyme immobilization. RSC

Citation: Yoshida N, Suzuki K, Ito S (2026) Comparative Analysis of enzyme immobilization techniques versus Standard Care in low bioreactor yield. Arch Biochem Eng, 1(1): 01-06.

- Advances, 4(4): 1583-1600. DOI: [10.1039/C3RA45991H](https://doi.org/10.1039/C3RA45991H)
25. Zaak, H., Siar, E.H., Kornecki, J.F., Fernandez-Lopez, L., Pedrero, S.G., Virgen-Ortiz, J.J., Fernandez-Lafuente, R. (2017) Effect of immobilization rate, at a low ionic strength, on the properties of biocatalysts. Process Biochemistry, 56: 117-123. DOI: [10.1016/j.procbio.2017.02.024](https://doi.org/10.1016/j.procbio.2017.02.024)
26. Tufvesson, P., Lima-Ramos, J., Nordblad, M., Woodley, J.M. (2011) Guidelines and cost analysis for catalyst production in biocatalytic processes. Organic Process Research & Development, 15(1): 266-274. DOI: [10.1021/op1002165](https://doi.org/10.1021/op1002165)
27. Illanes, A., Cauerhff, A., Wilson, L., Castro, G.R. (2012) Recent trends in biocatalysis engineering. Bioresource Technology, 115: 48-57. DOI: [10.1016/j.biortech.2011.12.050](https://doi.org/10.1016/j.biortech.2011.12.050)
28. DiCosimo, R., McAuliffe, J., Poulouse, A.J., Bohlmann, G. (2013) Industrial use of immobilized enzymes. Chemical Society Reviews, 42(15): 6437-6474. DOI: [10.1039/c3cs35506c](https://doi.org/10.1039/c3cs35506c)
29. Straathof, A.J.J., Panke, S., Schmid, A. (2002) The production of fine chemicals by biotransformations. Current Opinion in Biotechnology, 13(6): 548-556. DOI: [10.1016/S0958-1669\(02\)00360-9](https://doi.org/10.1016/S0958-1669(02)00360-9)
30. Garcia-Ochoa, F., Gomez, E. (2009) Bioreactor scale-up and oxygen transfer rate in microbial processes: an overview. Biotechnology Advances, 27(2): 153-176. DOI: [10.1016/j.biotechadv.2008.10.006](https://doi.org/10.1016/j.biotechadv.2008.10.006)
31. Chen, R.R. (2007) Permeability issues in whole-cell bioprocesses and cellular membrane engineering. Applied Microbiology and Biotechnology, 74(4): 730-738. DOI: [10.1007/s00253-006-0811-x](https://doi.org/10.1007/s00253-006-0811-x)
32. Fernandez-Lafuente, R. (2009) Stabilization of multimeric enzymes: strategies to prevent subunit dissociation. Enzyme and Microbial Technology, 45(6-7): 405-418. DOI: [10.1016/j.enzmictec.2009.08.009](https://doi.org/10.1016/j.enzmictec.2009.08.009)
33. Bilal, M., Iqbal, H.M.N. (2019) Naturally-derived biopolymers: potential platforms for enzyme immobilization. International Journal of Biological Macromolecules, 130: 462-482. DOI: [10.1016/j.ijbiomac.2019.02.152](https://doi.org/10.1016/j.ijbiomac.2019.02.152)
34. Ge, J., Lei, J., Zare, R.N. (2012) Protein-inorganic hybrid nanoflowers. Nature Nanotechnology, 7(7): 428-432. DOI: [10.1038/nnano.2012.80](https://doi.org/10.1038/nnano.2012.80)