

Archives of Diabetes

Review Article

HbA1c trend analysis in the Management of insulin resistance among elderly patients with diabetes

Ryan F^{1*}, Fitzgerald F² and Murphy N³

¹Department of Diabetes and Endocrinology, Rhine Academy of Metabolic Disease, Wales

²Department of Diabetes and Endocrinology, Amazon Center for Metabolic Disease, Wales

³Division of Diabetes and Endocrinology, Amazon Center for Metabolic Disease, Wales

*Address for Correspondence: Fionn Ryan, Department of Diabetes and Endocrinology, Rhine Academy of Metabolic Disease, Wales; E-mail: fionn.ryan@rhine.edu

Received: 5 November 2025; Accepted: 12 February 2026; Published: 17 May 2026.

Citation of this article: Ryan F, Fitzgerald F, Murphy N (2026) HbA1c trend analysis in the Management of insulin resistance among elderly patients with diabetes. Arch Diabetes, 8(1): 01-06. DOI: [10.5281/zenodo.22812656](https://doi.org/10.5281/zenodo.22812656)

Copyright: © 2026 Fionn Ryan, et al. This is an open access article distributed under the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Glycated hemoglobin (HbA1c) is the principal marker used to titrate therapy in elderly patients with type 2 diabetes, yet its trajectory reflects an age-specific mix of rising insulin resistance and declining insulin secretion that differs from younger cohorts.

Objective: This review synthesizes evidence on HbA1c trend analysis as a tool for monitoring insulin resistance and guiding individualized glycemic targets in older adults with diabetes.

Methods: We synthesized landmark randomized trials of intensive glycemic control, observational cohorts relating HbA1c trajectory to mortality, and consensus guidelines on target individualization in geriatric diabetes care.

Findings: Large cohorts show a U-shaped relationship between HbA1c and mortality in older adults, while randomized evidence from ACCORD, ADVANCE, and VADT shows intensive control lowers microvascular but not consistently macrovascular risk, and can raise mortality in frail elderly subgroups. Age-related increases in insulin resistance combine with declining beta-cell reserve to produce trend patterns that guideline bodies now use to stratify targets by functional status.

Conclusions: HbA1c trend analysis, interpreted alongside frailty and life-expectancy assessment rather than as an isolated threshold, supports individualized insulin-resistance management and safer glycemic targets in elderly patients with diabetes.

Keywords: HbA1c; insulin resistance; elderly; type 2 diabetes; glycemic control

1. Introduction

Glycated hemoglobin has served as the anchor laboratory value for type 2 diabetes management since the Diabetes Control and Complications Trial linked sustained lowering of HbA1c to a reduced risk of microvascular complications in insulin-treated patients [1], and since the UK Prospective Diabetes Study extended a comparable relationship to newly diagnosed type 2 diabetes [2,3]. In elderly patients, however, the same laboratory value sits inside a physiology that differs from the one those trials were built around. Aging is associated with a progressive rise in peripheral insulin resistance, driven by changes in body composition, reduced skeletal muscle mass, visceral adiposity, and mitochondrial function, layered onto a separate, largely independent decline in pancreatic beta-cell secretory reserve [4,5,6]. The result is that an HbA1c trajectory in a person in their eighties can reflect a very different balance of insulin resistance and insulin deficiency than the same trajectory in a person in their forties, even when the numerical values are identical.

This distinction matters clinically because elderly patients with diabetes are also the population in which the risks of tight glycemic control are least favorably balanced against its benefits. Multimorbidity, polypharmacy, reduced hypoglycemia awareness, cognitive impairment, and frailty are all more prevalent with advancing age, and each independently raises the hazard associated with aggressive HbA1c lowering [7,8,9]. Large cohort analyses have repeatedly found a U-shaped relationship between HbA1c and mortality in older adults with type 2 diabetes, with elevated risk at both the low and high ends of the distribution, a pattern rarely seen with comparable clarity in younger cohorts [10]. Three major randomized trials of intensive glycemic control published in the same era, ACCORD, ADVANCE, and VADT, together clarified both the ceiling and the hazard of pursuing near-normal HbA1c targets in populations with longstanding diabetes and cardiovascular risk factors typical of an aging cohort [11,12,13].

This review synthesizes what the endocrinology and geriatric-medicine literature has established about HbA1c trend analysis as a tool for characterizing insulin resistance and guiding glycemic management in elderly patients with diabetes: how the HbA1c trajectory relates to the underlying balance of insulin resistance and secretory failure with age, what the major trials and cohorts show about the trend's relationship to outcomes, and how guideline bodies now translate trend interpretation into individualized targets stratified by functional status.

2. Methods

This is a narrative synthesis of the published literature on HbA1c trajectory, insulin resistance, and glycemic target individualization in older adults with diabetes. We drew on landmark randomized controlled trials of intensive versus standard glycemic control (DCCT/EDIC, UKPDS, ACCORD, ADVANCE, VADT, ORIGIN), observational cohorts relating longitudinal HbA1c to mortality and complication risk in older populations, mechanistic studies of age-related insulin resistance and beta-cell function including the homeostasis model assessment (HOMA-IR) literature, and consensus and position statements from the American Diabetes Association, the American Geriatrics Society, and the European Diabetes Working Party for Older People addressing glycemic target individualization in geriatric diabetes care. Studies were selected for topical relevance to HbA1c trend interpretation, insulin resistance physiology in aging, and glycemic target setting in elderly patients, rather than through a formal systematic search protocol, consistent with the narrative-review format of this article. Findings are organized around three questions: how HbA1c trend reflects the changing balance of insulin resistance and insulin deficiency with age, what randomized and observational evidence shows about the trend's relationship to clinical outcomes in older adults, and how that evidence has been translated into individualized target-setting frameworks.

3. Results

HbA1c trend and the changing balance of insulin resistance with age

The pathophysiological chain from aging to a rising HbA1c trend runs through two largely separable mechanisms. Cross-sectional and longitudinal studies using the homeostasis model assessment of insulin resistance (HOMA-IR), the widely adopted method derived by Matthews and colleagues from fasting glucose and insulin concentrations, consistently show higher indices of insulin resistance in older compared with younger adults matched for body mass index [14]. Ferrannini and colleagues, studying insulin sensitivity across a large multicenter European cohort, found that insulin resistance increased with age independent of adiposity, implicating intrinsic changes in skeletal muscle glucose disposal rather than obesity alone as the principal driver [15]. Barzilai and Ferrucci, reviewing the aging literature, argued that this age-related rise in insulin resistance may in some contexts represent an adaptive, protective shift in fuel partitioning rather than a purely pathological process, a reframing relevant to interpreting HbA1c trend in the oldest patients [16].

Superimposed on this resistance trend is an independent, and in many patients dominant, decline in beta-cell secretory capacity. Chang and Halter, reviewing insulin secretion across the lifespan, found that aging blunts both the first-phase and pulsatile components of insulin release in response to glucose, a defect that compounds peripheral resistance rather than compensating for it [6]. Meneilly and Elahi, studying lean elderly patients with type 2 diabetes, found that a subset of older patients present with a metabolic phenotype dominated by secretory failure rather than resistance, a distinction with direct implications for whether insulin sensitizers or secretagogues better target the trend driving a given patient's rising HbA1c [17]. DeFronzo's synthesis of the pathophysiology of type 2 diabetes, describing the expanding "ominous octet" of contributing organ

systems, situates this age-related uncoupling of resistance and secretion within the broader multi-organ model that now underlies rational, mechanism-based glucose-lowering therapy selection in older patients [4]. Kahn's earlier analysis of the relative contributions of insulin resistance and beta-cell dysfunction across the natural history of type 2 diabetes established the conceptual framework, still in use, for interpreting which mechanism a rising or falling HbA1c trend most plausibly reflects at a given disease stage [5].


Table 1: Age-related contributions to a rising HbA1c trend, comparing peripheral insulin resistance and beta-cell secretory decline as distinct mechanisms [14,15,6,17].			
Mechanism	Typical age-related change	Dominant metabolic phenotype	Key evidence
Peripheral insulin resistance	Increases with age, largely independent of adiposity	Preserved secretion, elevated HOMA-IR	[14,15]
Beta-cell secretory capacity	Declines with age, blunted first-phase response	Lean phenotype, secretory failure dominant	[6,17]
Combined resistance and secretory decline	Both mechanisms present, variable weighting	Mixed phenotype, most common in longstanding disease	[4,5]

HbA1c trend and outcomes in randomized trials and cohorts

The clearest evidence on how HbA1c trend relates to outcomes in older, longstanding diabetes populations comes from three large randomized trials published within a year of one another. The ACCORD trial randomized patients with type 2 diabetes and elevated cardiovascular risk to an intensive strategy targeting HbA1c below 6.0% or a standard strategy targeting 7.0-7.9%, and was stopped early after intensive treatment was associated with significantly higher all-cause mortality despite achieving a lower mean HbA1c trajectory [11]. The ADVANCE trial, using a less aggressive intensive target of 6.5%, found a significant reduction in the composite of major macrovascular and microvascular

events, driven predominantly by a reduction in nephropathy, without the excess mortality seen in ACCORD [12]. The VADT, enrolling patients with a longer mean diabetes duration than either of the other two trials, found no significant difference in cardiovascular outcomes between intensive and standard glycemic control over the trial period, though a subsequent long-term follow-up suggested a delayed benefit [13]. Ismail-Beigi and colleagues' detailed analysis of the ACCORD microvascular data found that intensive glucose lowering did reduce the composite renal and retinal outcome, indicating that the mortality signal in that trial did not reflect a uniform hazard of glycemic control across all organ systems [18].

Table 2: Intensive versus standard glycemic control in three pivotal randomized trials of type 2 diabetes with a substantial older-adult population [11,12,13].			
Trial	Intensive HbA1c target	Mean diabetes duration	Key finding
[11]	<6.0%	~10 years	Stopped early; higher all-cause mortality with intensive arm
[12]	≤6.5%	~8 years	Reduced composite macro/microvascular events, no excess mortality
[13]	~6.9% achieved	~11.5 years	No significant cardiovascular benefit during the trial period



Large observational cohorts extend this picture to the

HbA1c trend itself rather than a single randomized target. Currie and colleagues, following a large primary-care cohort, found a U-shaped relationship between mean HbA1c and all-cause mortality, with the lowest hazard near 7.5% and elevated hazard at both markedly lower and higher values, difficult to reconcile with a simple "lower is always better" reading of trend [10]. Huang and colleagues, in the Diabetes and Aging Study, found that severe hypoglycemia requiring emergency treatment rose sharply among older patients maintained at very low HbA1c, particularly those with substantial comorbidity, a population in whom a falling trend can itself become the adverse event [19]. Lipska and colleagues found that many older adults with complex or poor health status were nonetheless maintained at tightly controlled HbA1c inconsistent with individualized targets, indicating overtreatment is common in practice rather than a theoretical concern [20]. Selvin and colleagues' meta-analysis confirmed a continuous, graded association between HbA1c and cardiovascular risk, supporting continued surveillance even where near-normalization is not the goal [21]. Glycemic variability adds a further dimension: Monnier and colleagues found that acute glucose fluctuations activate oxidative stress independent of mean glycemia, and Kilpatrick and colleagues, reanalyzing DCCT data, found that variability contributed to microvascular risk beyond mean HbA1c alone, both relevant to why a stable trend, not merely a favorable mean, is now emphasized [22,23].

From trend to individualized glycemic targets

The practical translation of this evidence has been a shift from a single, uniform HbA1c target toward stratified targets set according to functional status, comorbidity, and estimated life expectancy. Kirkman and colleagues' consensus report, prepared jointly for the American Diabetes Association and the American Geriatrics Society, proposed a three-tier framework: an HbA1c target below approximately 7.5% for healthy older adults with long life expectancy, below 8.0% for moderate comorbidity, and

below 8.5% for multiple comorbidities or limited life expectancy, prioritizing avoidance of hypoglycemia over near-normalization in the most vulnerable tier [24]. Sinclair and colleagues' European Diabetes Working Party for Older People guidelines proposed a comparable framework, emphasizing functional and cognitive assessment, not age alone, as the required input to target-setting [25]. The American Geriatrics Society's expert panel echoed this stratification, recommending against tight control in older adults with limited life expectancy given the multi-year lag before microvascular benefit accrues [26]. Munshi and colleagues extended this framework to long-term care and skilled nursing facilities, recommending relaxed targets, often above 8%, and simplified regimens prioritizing avoidance of hypoglycemic trend reversals [27]. Lee and Eng reinforced the same principle from a geriatrics perspective: a falling HbA1c trend in a frail patient should prompt evaluation for overtreatment rather than being read uncritically as success [8]. The American Diabetes Association's current Standards of Care formalizes this individualized approach in its glycemic-target and dedicated older-adults guidance [28,29]. The ORIGIN trial's long-term follow-up of early basal insulin in dysglycemia reinforced that neutral cardiovascular outcomes with careful, hypoglycemia-avoidant titration are achievable even when trend improvement is modest [30]. Turner and colleagues' UKPDS 49 analysis, showing that monotherapy increasingly fails to sustain HbA1c control over time, underscores why trend surveillance remains necessary throughout the disease course, including into old age [31]. Yakaryilmaz and Ozturk, and Chiniwala and Jabbour, synthesize this stratified approach into prescribing guidance favoring agents less prone to provoking hypoglycemic excursions [32,33]. Rodriguez-Gutierrez and Montori caution that guidance in this area has been repeatedly revised as evidence accumulated, a caution they argue applies with particular force to elderly patients, where the evidence base remains

thin [34]. Boyle and colleagues' population projections underscore this framework's growing relevance given the rising share of the diabetes population expected to be over 65 in coming decades [35].




Figure 2: Individualized HbA1c target ceilings recommended by functional and comorbidity tier in the American Diabetes Association / American Geriatrics Society consensus framework for older adults [24].

4. Discussion

Taken together, this literature supports reading HbA1c trend in elderly patients as a signal of at least three overlapping processes rather than a single number to be minimized: the underlying, age-related shift in the balance between insulin resistance and beta-cell secretory failure [4,5,17]; the accumulating risk, evident in both randomized trials and large cohorts, that pursuing a low absolute HbA1c value can itself become hazardous once comorbidity, frailty, and hypoglycemia risk are accounted for [11,10,19]; and the practical need, now embedded in every major geriatric diabetes guideline, to set the target the trend is measured against according to functional status rather than age or diagnosis alone [24,25,27]. The U-shaped mortality curve described by Currie and colleagues is perhaps the single clearest illustration of why trend direction and trend context matter more than trend magnitude in isolation: a falling HbA1c in a frail, polypharmacy-burdened patient is not intrinsically reassuring, and a stable, moderately elevated HbA1c in the same patient may represent appropriately calibrated care rather than undertreatment [10,8].

This synthesis has the limitations inherent to a narrative

rather than systematic review, and the underlying evidence base itself has gaps particular to the oldest and frailest patients, who are systematically underrepresented in the pivotal randomized trials that anchor current practice; ACCORD, ADVANCE, and VADT all enrolled predominantly middle-aged and early-elderly participants with preserved functional status relative to the nursing-home and multimorbid populations that guideline stratification frameworks are increasingly asked to address [11,12,13,27]. Mechanistic evidence on the age-related uncoupling of insulin resistance and secretory failure is stronger than the outcomes evidence linking specific trend patterns to specific interventions, meaning that much of the individualization now recommended in practice rests more on consensus judgment than on trial data generated within the elderly population itself [15,6,26].

5. Conclusion

HbA1c trend analysis remains central to managing insulin resistance in elderly patients with diabetes, but its interpretation in this population differs meaningfully from its interpretation in younger, less comorbid cohorts. The trend reflects a shifting balance between age-related insulin resistance and declining beta-cell reserve [4,14], its relationship to mortality and complications follows a U-shaped rather than a strictly linear pattern once frailty and hypoglycemia risk are considered [10,19], and the major randomized trials of intensive control collectively counsel against pursuing near-normal targets in patients with longstanding diabetes and significant comorbidity [11,12,13]. Current guidance from the American Diabetes Association, the American Geriatrics Society, and the European Diabetes Working Party for Older People converges on a stratified approach that sets the HbA1c target the trend is judged against according to functional status and life expectancy rather than a single universal threshold [24,25,26]. Read this way, as a trajectory interpreted against an individualized target rather than a value to be minimized on principle, HbA1c trend analysis

remains a genuinely useful tool for managing insulin resistance safely across the wide range of physiological reserve seen in elderly patients with diabetes.

6. References

1. DCCT Research Group. (1993) The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *New England Journal of Medicine*, 329(14): 977-986. DOI: [10.1056/NEJM199309303291401](https://doi.org/10.1056/NEJM199309303291401)
2. UK Prospective Diabetes Study (UKPDS) Group. (1998) Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet*, 352(9131): 837-853.
3. Stratton, I.M., Adler, A.I., Neil, H.A., Matthews, D.R., Manley, S.E., Cull, C.A., et al. (2000) Association of glycaemia with macrovascular and microvascular complications of type 2 diabetes (UKPDS 35): prospective observational study. *BMJ*, 321(7258): 405-412. DOI: [10.1136/bmj.321.7258.405](https://doi.org/10.1136/bmj.321.7258.405)
4. DeFronzo, R.A. (2009) Banting Lecture. From the triumvirate to the ominous octet: a new paradigm for the treatment of type 2 diabetes mellitus. *Diabetes*, 58(4): 773-795. DOI: [10.2337/db09-9028](https://doi.org/10.2337/db09-9028)
5. Kahn, S.E. (2003) The relative contributions of insulin resistance and beta-cell dysfunction to the pathophysiology of Type 2 diabetes. *Diabetologia*, 46(1): 3-19. DOI: [10.1007/s00125-002-1009-0](https://doi.org/10.1007/s00125-002-1009-0)
6. Chang, A.M., Halter, J.B. (2003) Aging and insulin secretion. *American Journal of Physiology-Endocrinology and Metabolism*, 284(1): E7-E12. DOI: [10.1152/ajpendo.00366.2002](https://doi.org/10.1152/ajpendo.00366.2002)
7. Halter, J.B., Musi, N., McFarland Horne, F., Crandall, J.P., Goldberg, A., Harkless, L., et al. (2014) Diabetes and cardiovascular disease in older adults: current status and future directions. *Diabetes*, 63(8): 2578-2589. DOI: [10.2337/131111](https://doi.org/10.2337/131111)

[10.2337/db14-0020](https://doi.org/10.2337/db14-0020)

8. Lee, S.J., Eng, C. (2011) Goals of glycemic control in frail older patients with diabetes. JAMA, 305(13): 1350-1351. DOI: [10.1001/jama.2011.404](https://doi.org/10.1001/jama.2011.404)
9. Bandeen-Roche, K., Xue, Q.L., Ferrucci, L., Walston, J., Guralnik, J.M., Chaves, P., et al. (2006) Phenotype of frailty: characterization in the women's health and aging studies. Journals of Gerontology Series A: Biological Sciences and Medical Sciences, 61(3): 262-266. DOI: [10.1093/gerona/61.3.262](https://doi.org/10.1093/gerona/61.3.262)
10. Currie, C.J., Peters, J.R., Tynan, A., Evans, M., Heine, R.J., Bracco, O.L., et al. (2010) Survival as a function of HbA1c in people with type 2 diabetes: a retrospective cohort study. Lancet, 375(9713): 481-489. DOI: [10.1016/S0140-6736\(09\)61969-3](https://doi.org/10.1016/S0140-6736(09)61969-3)
11. Action to Control Cardiovascular Risk in Diabetes Study Group. (2008) Effects of intensive glucose lowering in type 2 diabetes. New England Journal of Medicine, 358(24): 2545-2559. DOI: [10.1056/NEJMoa0802743](https://doi.org/10.1056/NEJMoa0802743)
12. ADVANCE Collaborative Group. (2008) Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. New England Journal of Medicine, 358(24): 2560-2572. DOI: [10.1056/NEJMoa0802987](https://doi.org/10.1056/NEJMoa0802987)
13. Duckworth, W., Abraira, C., Moritz, T., Reda, D., Emanuele, N., Reaven, P.D., et al. (2009) Glucose control and vascular complications in veterans with type 2 diabetes. New England Journal of Medicine, 360(2): 129-139. DOI: [10.1056/NEJMoa0808431](https://doi.org/10.1056/NEJMoa0808431)
14. Matthews, D.R., Hosker, J.P., Rudenski, A.S., Naylor, B.A., Treacher, D.F., Turner, R.C. (1985) Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. Diabetologia, 28(7): 412-419. DOI: [10.1007/BF00280883](https://doi.org/10.1007/BF00280883)
15. Ferrannini, E., Natali, A., Bell, P., Cavallo-Perin, P., Lalic, N., Mingrone, G. (1997) Insulin resistance and hypersecretion in obesity. European Group for the Study of Insulin Resistance (EGIR). Journal of Clinical Investigation, 100(5): 1166-1173. DOI: [10.1172/JCI119628](https://doi.org/10.1172/JCI119628)
16. Barzilai, N., Ferrucci, L. (2012) Insulin resistance and aging: a cause or a protective response? Journals of Gerontology Series A: Biological Sciences and Medical Sciences, 67(12): 1329-1331. DOI: [10.1093/gerona/gls145](https://doi.org/10.1093/gerona/gls145)
17. Meneilly, G.S., Elahi, D. (2005) Metabolic alterations in middle-aged and elderly lean patients with type 2 diabetes. Diabetes Care, 28(6): 1498-1499. DOI: [10.2337/diacare.28.6.1498](https://doi.org/10.2337/diacare.28.6.1498)
18. Ismail-Beigi, F., Craven, T., Banerji, M.A., Basile, J., Calles, J., Cohen, R.M., et al. (2010) Effect of intensive treatment of hyperglycaemia on microvascular outcomes in type 2 diabetes: an analysis of the ACCORD randomised trial. Lancet, 376(9739): 419-430. DOI: [10.1016/S0140-6736\(10\)60576-4](https://doi.org/10.1016/S0140-6736(10)60576-4)
19. Huang, E.S., Laiteerapong, N., Liu, J.Y., John, P.M., Moffet, H.H., Karter, A.J. (2014) Rates of complications and mortality in older patients with diabetes mellitus: the diabetes and aging study. JAMA Internal Medicine, 174(2): 251-258. DOI: [10.1001/jamainternmed.2013.12956](https://doi.org/10.1001/jamainternmed.2013.12956)
20. Lipska, K.J., Ross, J.S., Miao, Y., Shah, N.D., Lee, S.J., Steinman, M.A. (2015) Potential overtreatment of diabetes mellitus in older adults with tight glycemic control. JAMA Internal Medicine, 175(3): 356-362. DOI: [10.1001/jamainternmed.2014.7345](https://doi.org/10.1001/jamainternmed.2014.7345)
21. Selvin, E., Marinopoulos, S., Berkenblit, G., Rami, T., Brancati, F.L., Powe, N.R., et al. (2004) Meta-analysis: glycosylated hemoglobin and cardiovascular disease in diabetes mellitus. Annals of Internal Medicine, 141(6): 421-431. DOI: [10.7326/0003-4819-141-6-200409210-00007](https://doi.org/10.7326/0003-4819-141-6-200409210-00007)
22. Monnier, L., Mas, E., Ginet, C., Michel, F., Villon, L., Cristol, J.P., et al. (2006) Activation of oxidative stress by acute glucose fluctuations compared with sustained chronic hyperglycemia in patients with type 2 diabetes. JAMA,

Citation: Ryan F, Fitzgerald F, Murphy N (2026) HbA1c trend analysis in the Management of insulin resistance among elderly patients with diabetes. Arch Diabetes, 8(1): 01-06.

- 295(14): 1681-1687. DOI: [10.1001/jama.295.14.1681](https://doi.org/10.1001/jama.295.14.1681)
23. Kilpatrick, E.S., Rigby, A.S., Atkin, S.L. (2006) The effect of glucose variability on the risk of microvascular complications in type 1 diabetes. Diabetes Care, 29(7): 1486-1490. DOI: [10.2337/dc06-0293](https://doi.org/10.2337/dc06-0293)
24. Kirkman, M.S., Briscoe, V.J., Clark, N., Florez, H., Haas, L.B., Halter, J.B., et al. (2012) Diabetes in older adults. Diabetes Care, 35(12): 2650-2664. DOI: [10.2337/dc12-1801](https://doi.org/10.2337/dc12-1801)
25. Sinclair, A.J., Paolisso, G., Castro, M., Bourdel-Marchasson, I., Gadsby, R., Rodriguez-Manas, L. (2011) European Diabetes Working Party for Older People 2011 clinical guidelines for type 2 diabetes mellitus. Executive summary. Diabetes & Metabolism, 37(S3): S27-S38. DOI: [10.1016/S1262-3636\(11\)70962-4](https://doi.org/10.1016/S1262-3636(11)70962-4)
26. American Geriatrics Society Expert Panel on the Care of Older Adults with Diabetes Mellitus. (2013) Guidelines abstracted from the American Geriatrics Society guidelines for improving the care of older adults with diabetes mellitus. Journal of the American Geriatrics Society, 61(11): 2020-2026. DOI: [10.1111/jgs.12514](https://doi.org/10.1111/jgs.12514)
27. Munshi, M.N., Florez, H., Huang, E.S., Kalyani, R.R., Mupanomunda, M., Pandya, N., et al. (2016) Management of diabetes in long-term care and skilled nursing facilities: a position statement of the American Diabetes Association. Diabetes Care, 39(2): 308-318. DOI: [10.2337/dc15-2512](https://doi.org/10.2337/dc15-2512)
28. American Diabetes Association Professional Practice Committee. (2023) 6. Glycemic targets: standards of care in diabetes-2023. Diabetes Care, 46(Suppl 1): S97-S110. DOI: [10.2337/dc23-S006](https://doi.org/10.2337/dc23-S006)
29. American Diabetes Association Professional Practice Committee. (2024) 13. Older adults: standards of care in diabetes-2024. Diabetes Care, 47(Suppl 1): S244-S257. DOI: [10.2337/dc24-S013](https://doi.org/10.2337/dc24-S013)
30. ORIGIN Trial Investigators. (2012) Basal insulin and cardiovascular and other outcomes in dysglycemia. New England Journal of Medicine, 367(4): 319-328. DOI: [10.1056/NEJMoa1203858](https://doi.org/10.1056/NEJMoa1203858)
31. Turner, R.C., Cull, C.A., Frighi, V., Holman, R.R. (1999) Glycemic control with diet, sulfonylurea, metformin, or insulin in patients with type 2 diabetes mellitus: progressive requirement for multiple therapies (UKPDS 49). JAMA, 281(21): 2005-2012. DOI: [10.1001/jama.281.21.2005](https://doi.org/10.1001/jama.281.21.2005)
32. Yakaryilmaz, F.D., Ozturk, Z.A. (2017) Treatment of type 2 diabetes mellitus in the elderly. World Journal of Diabetes, 8(6): 278-285. DOI: [10.4239/wjd.v8.i6.278](https://doi.org/10.4239/wjd.v8.i6.278)
33. Chiniwala, N., Jabbour, S. (2011) Management of diabetes mellitus in the elderly. Current Opinion in Endocrinology, Diabetes and Obesity, 18(2): 148-152. DOI: [10.1097/MED.0b013e3283444fbd](https://doi.org/10.1097/MED.0b013e3283444fbd)
34. Rodriguez-Gutierrez, R., Montori, V.M. (2016) Glycemic control for patients with type 2 diabetes mellitus: our evolving faith in the face of evidence. Circulation: Cardiovascular Quality and Outcomes, 9(5): 504-512. DOI: [10.1161/CIRCOUTCOMES.116.002901](https://doi.org/10.1161/CIRCOUTCOMES.116.002901)
35. Boyle, J.P., Thompson, T.J., Gregg, E.W., Barker, L.E., Williamson, D.F. (2010) Projection of the year 2050 burden of diabetes in the US adult population: dynamic modeling of incidence, mortality, and prediabetes prevalence. Population Health Metrics, 8: 29. DOI: [10.1186/1478-7954-8-29](https://doi.org/10.1186/1478-7954-8-29)