

# Archives of Diabetes

## Research Article

# Comparative Analysis of structured lifestyle intervention versus Standard Care in gestational diabetes

Walsh N<sup>1\*</sup> , Kelly S<sup>2</sup>  and Fitzgerald O<sup>3</sup> 

<sup>1</sup>Department of Diabetes and Endocrinology, Clyde Research Center for Metabolic Disease, Wales

<sup>2</sup>Division of Diabetes and Endocrinology, Parana Institute for Advanced Metabolic Disease, Canada

<sup>3</sup>Division of Diabetes and Endocrinology, Fraser Institute of Metabolic Disease, Ireland

\*Address for Correspondence: Niamh Walsh, Department of Diabetes and Endocrinology, Clyde Research Center for Metabolic Disease, Wales; E-mail: [niamh.walsh@clyde.edu](mailto:niamh.walsh@clyde.edu)

Received: 21 March 2026; Accepted: 27 June 2026; Published: 16 September 2026.

Citation of this article: Walsh N, Kelly S, Fitzgerald O (2026) Comparative Analysis of structured lifestyle intervention versus Standard Care in gestational diabetes. Arch Diabetes, 8(1): 01-06. DOI: [10.5281/zenodo.22812634](https://doi.org/10.5281/zenodo.22812634)

Copyright: © 2026 Niamh Walsh, et al. This is an open access article distributed under the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

**Background:** Gestational diabetes mellitus (GDM) affects a substantial proportion of pregnancies worldwide and raises the risk of macrosomia, caesarean delivery, and later maternal type 2 diabetes. Structured lifestyle intervention programmes combining individualised dietary counselling, supervised physical activity, and structured glucose monitoring have increasingly been compared against standard antenatal care for glycaemic and obstetric outcomes.

**Objective:** This comparative analysis synthesizes randomized and quasi-randomized evidence contrasting structured lifestyle intervention with standard care in women with or at risk of GDM, focusing on maternal glycaemic control, gestational weight gain, and neonatal outcomes.

**Methods:** Trials, systematic reviews, and meta-analyses addressing structured dietary, exercise, and combined lifestyle programmes in GDM were synthesized narratively, with outcome patterns compared descriptively across structured intervention and standard-care arms.

**Results:** Structured lifestyle intervention was consistently associated with lower rates of macrosomia and large-for-gestational-age infants, more favourable gestational weight gain, and modestly improved glycaemic control relative to standard care, with less consistent effects on caesarean delivery and preeclampsia.

**Conclusion:** Structured lifestyle intervention offers measurable advantages over standard care for glycaemic and selected neonatal outcomes in GDM, supporting its integration into routine antenatal management alongside individualised risk stratification.

**Keywords:** gestational diabetes; lifestyle intervention; standard care; glycemic control; pregnancy outcomes

## 1. Introduction

Gestational diabetes mellitus (GDM) is one of the most common metabolic complications of pregnancy, and its global prevalence has risen alongside increasing rates of maternal obesity and advancing maternal age at conception [1]. The Hyperglycemia and Adverse Pregnancy Outcomes (HAPO) study established a continuous, graded relationship between maternal glucose concentrations below the threshold for overt diabetes and adverse outcomes including birth weight above the 90th percentile, primary caesarean delivery, and neonatal hypoglycaemia, a finding that reshaped diagnostic thresholds worldwide [2]. The International Association of Diabetes and Pregnancy Study Groups (IADPSG) subsequently issued consensus criteria that substantially lowered the glycaemic threshold for diagnosis, increasing the proportion of pregnancies classified as GDM in many populations and intensifying the clinical question of which management strategy best improves outcomes once the diagnosis is made [3,4].

Standard antenatal care for GDM has historically consisted of general dietary advice at diagnosis, intermittent self-monitoring of blood glucose, and pharmacological therapy reserved for women who fail glycaemic targets, with little structured reinforcement of behaviour change [5]. Structured lifestyle intervention, by contrast, bundles individualised dietary counselling from a qualified dietitian, a prescribed and supervised physical activity component, structured glucose monitoring with feedback, and scheduled follow-up designed to sustain adherence [6]. Two landmark randomized trials, ACHOIS and the US Maternal-Fetal Medicine Units Network mild GDM trial, demonstrated that active management of even mild hyperglycaemia reduces serious perinatal morbidity and macrosomia compared with usual care, establishing that structured management, not diagnosis alone, drives outcome improvement [7,8].

A substantial literature has examined whether more

intensively structured lifestyle programmes confer additional benefit once GDM is diagnosed, and a parallel literature has examined whether structured intervention initiated earlier in pregnancy can prevent GDM in high-risk women [9,10,11]. These two related but distinct questions, treatment of established GDM and primary prevention, are often conflated in narrative summaries; this analysis addresses both explicitly, synthesizing comparative evidence across maternal glycaemic, metabolic, and neonatal domains to characterise where the advantage of structured intervention is most consistent.

## 2. Methods

We synthesized randomized controlled trials, Cochrane systematic reviews, and meta-analyses that compared a structured lifestyle intervention arm, defined as a programme combining individualised dietary counselling, a prescribed physical activity component, and structured monitoring or follow-up, against a standard-care or usual-care comparator in women with diagnosed GDM or at elevated risk of developing it. Sources were drawn from systematic reviews addressing dietary intervention [12,13], exercise-based intervention [10,14], combined lifestyle prevention [9,15,16], and treatment of established GDM [17,18,19], supplemented by primary trials frequently cited within those reviews [20,21,22,23,24,25,26]. Outcome domains extracted were maternal glycaemic control, gestational weight gain, GDM incidence in prevention trials, mode of delivery, and neonatal outcomes including macrosomia, large-for-gestational-age (LGA) birth weight, and neonatal hypoglycaemia. Because studies varied in population, intervention intensity, and outcome definitions, findings were synthesized narratively rather than pooled quantitatively. Current ACOG and ADA guidance on GDM management was used to frame clinical context throughout [27,28].

## 3. Comparative maternal glycaemic outcomes

Across dietary and combined lifestyle trials, structured

intervention has been associated with modestly but consistently better glycaemic control than standard advice-only care. Yamamoto and colleagues, in a systematic review and meta-analysis of randomized trials examining modified dietary interventions in established GDM, found that structured low-glycaemic-index and similarly modified diets reduced the need for insulin therapy and improved fasting and postprandial glucose relative to standard dietary advice, without compromising neonatal birth weight [12]. Viana and colleagues reached a broadly concordant conclusion in their meta-analysis of dietary randomized trials, reporting that structured low-glycaemic-index dietary counselling reduced insulin requirement and improved glycaemic outcomes compared with conventional dietary advice, while higher-fibre and energy-restricted structured diets showed more heterogeneous effects [13].

**Table 1: Overview of major comparative trials of structured lifestyle intervention versus standard care in gestational diabetes mellitus prevention and treatment [7,8,20,21,22].**

| Trial          | Population                | Structured intervention                               | Comparator              |
|----------------|---------------------------|-------------------------------------------------------|-------------------------|
| ACHOIS         | Established GDM           | Dietary advice, glucose monitoring, insulin as needed | Routine antenatal care  |
| Mild GDM trial | Mild GDM                  | Structured diet and monitoring protocol               | Usual prenatal care     |
| RADIEL         | History of GDM or obesity | Dietitian-led diet and activity counselling           | Standard antenatal care |
| UPBEAT         | Obese pregnant women      | Structured behavioural diet and activity coaching     | Standard antenatal care |
| LIMIT          | Overweight/obese women    | Structured dietary and lifestyle advice               | Standard antenatal care |

The Cochrane review by Brown and colleagues, synthesizing lifestyle interventions for women already diagnosed with GDM, concluded that structured dietary and lifestyle advice was associated with reduced macrosomia and improved glycaemic control compared with standard care, though certainty of evidence varied

across outcomes [17]. Horvath and colleagues similarly found that active glycaemic management strategies incorporating structured dietary and monitoring components reduced macrosomia and shoulder dystocia relative to usual-care management [19]. Wang and colleagues, in a randomized trial of supervised exercise in overweight and obese women, reported improved glucose tolerance and a lower GDM incidence in the exercise arm relative to standard care [14]. The DALI project, a multicentre lifestyle trial across European centres, similarly linked greater achieved physical activity within the structured arm to better glycaemic control [26].

**4. Comparative gestational weight gain and metabolic outcomes**

Gestational weight gain is one of the most consistently favourable outcomes associated with structured lifestyle intervention. The UPBEAT trial, a large multicentre trial of structured behavioural diet and activity coaching in obese pregnant women, reduced gestational weight gain and improved maternal dietary indices relative to standard care, though it did not significantly reduce the composite outcome of GDM and LGA birth [21]. The LIMIT trial likewise found that structured dietary and lifestyle counselling reduced excessive gestational weight gain compared with standard care, without significantly reducing LGA birth as a primary outcome [22].

Prevention-focused trials in women at elevated risk report similar patterns. The Finnish RADIEL study randomized women with a history of GDM or obesity to dietitian-led structured counselling versus standard antenatal care and found roughly one-third lower GDM incidence with structured intervention, alongside more favourable weight gain [20]. Luoto and colleagues, in a Finnish cluster-randomized trial of primary-care-delivered counselling, reported a reduced combined incidence of GDM and LGA birth with structured counselling, particularly among overweight women [23]. The St Carlos study randomized women to a Mediterranean-diet

structured intervention, supplemented with olive oil and nuts, versus standard advice and found substantially lower GDM incidence in the intervention group [25]. The BAMBINO feasibility trial found structured exercise intervention in obese women achievable and associated with more favourable weight gain [24]. Bao and colleagues showed that greater postpartum activity among women with prior GDM was associated with lower risk of progression to type 2 diabetes, suggesting the metabolic benefit of structured activity extends beyond pregnancy [29].




**Figure 1:** Incidence of gestational diabetes mellitus among women receiving structured lifestyle intervention versus standard care in two prevention trials, the Finnish RADIEL study and the Madrid St Carlos Mediterranean-diet study, both showing lower incidence with structured intervention [20,25].

Meta-analytic syntheses support a genuine, if modest, aggregate effect. Song and colleagues, pooling prevention trials, reported a significant reduction in GDM incidence with structured diet-and-exercise counselling, most pronounced when intervention began earlier and was more intensive [15]. Guo and colleagues, using meta-regression, found effectiveness strongly tied to programme structure and adherence support rather than intervention type alone [16]. The Cochrane reviews of diet-and-exercise and exercise-only prevention reached compatible conclusions, with more heterogeneous effects for exercise alone and generally low-to-moderate certainty of evidence [9,10].

## 5. Comparative neonatal and perinatal outcomes

The obstetric and neonatal literature shows its clearest,

most consistent advantage in macrosomia and LGA birth weight. The Cochrane reviews of dietary advice for preventing GDM and of treatment in established GDM both reported reduced macrosomia with structured dietary counselling relative to standard care, an effect reproduced across independent meta-analyses [11,17]. The ACHOIS trial reported substantially lower serious perinatal complications, including death, shoulder dystocia, fracture, and nerve palsy, with structured dietary advice, monitoring, and insulin as needed, compared with routine care [7]. The mild GDM trial reported similarly lower macrosomia, LGA birth, shoulder dystocia, and preeclampsia with structured treatment compared with usual prenatal care [8].




**Figure 2:** Serious perinatal complications and macrosomia in the ACHOIS trial, comparing structured dietary advice, glucose monitoring, and insulin therapy as needed against routine antenatal care without structured glycaemic management [7].

Hartling and colleagues, in a review for the US Preventive Services Task Force, concluded that structured treatment protocols reduced macrosomia, shoulder dystocia, and preeclampsia relative to standard treatment, informing subsequent screening guidance [18]. Neonatal hypoglycaemia, a theoretical concern with tighter glycaemic control, was not consistently increased by structured intervention: both ACHOIS and the mild GDM trial reported similar hypoglycaemia rates between arms, suggesting the glycaemic benefit is not achieved at the expense of neonatal risk [7,8].

## 6. Comparative mode of delivery and maternal outcomes

| Table 2: Comparative consistency of structured lifestyle intervention effects versus standard care across outcome domains, drawn from treatment and prevention trials and systematic reviews [17,18,16,30]. |                                       |                           |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------|---------------------------|
| Outcome domain                                                                                                                                                                                              | Effect of structured intervention     | Consistency across trials |
| Maternal glycaemic control                                                                                                                                                                                  | Improved, reduced insulin requirement | Consistent                |
| Gestational weight gain                                                                                                                                                                                     | Reduced excessive weight gain         | Consistent                |
| Macrosomia / LGA birth weight                                                                                                                                                                               | Reduced incidence                     | Consistent                |
| Caesarean delivery                                                                                                                                                                                          | Variable, often not significant       | Inconsistent              |
| Preeclampsia                                                                                                                                                                                                | Modest reduction in some trials       | Inconsistent              |
| Neonatal hypoglycaemia                                                                                                                                                                                      | No significant increase               | Consistent                |

Effects on caesarean delivery and hypertensive disorders are less consistent than glycaemic or macrosomia outcomes. The mild GDM trial found a modest preeclampsia reduction with structured treatment, but no significant effect on caesarean delivery rate [8]. UPBEAT and LIMIT, both in obese women, found no significant reduction in caesarean delivery despite favourable weight-gain effects, indicating that mode-of-delivery outcomes are influenced by factors a pregnancy-limited lifestyle intervention does not fully modify [21,22]. The Cochrane review of GDM treatment likewise reported inconsistent effects on caesarean rate, in contrast to the more uniform macrosomia reduction [17]. Zhang and colleagues noted that the metabolic benefits of structured diet and activity are more directly linked to glycaemic and fetal growth pathways than to the obstetric determinants of delivery mode, a plausible explanation for this pattern [30].

7. Discussion

Taken together, the literature supports a nuanced conclusion: structured lifestyle intervention, whether as prevention in at-risk women or first-line treatment in established GDM, produces a consistent, clinically meaningful advantage over standard care for glycaemic control, gestational weight gain, and macrosomia or LGA birth weight, while its advantage for caesarean delivery, preeclampsia, and neonatal hypoglycaemia is smaller and

less uniform [17,18,19]. This pattern fits the underlying physiology: structured diet and activity act directly on maternal glucose flux and fetal growth, the pathway most proximate to macrosomia, whereas caesarean delivery and preeclampsia are determined by a wider set of factors a pregnancy-limited programme cannot fully modify [30,27].

The meta-regression findings of Guo and colleagues indicate that benefit is not intrinsic to counselling in the abstract but depends on programme structure, adherence support, and timing [16]. This reconciles apparently discordant results: UPBEAT and LIMIT, large trials in high-risk obese populations, did not reduce their composite neonatal outcomes despite gains in weight gain and diet quality, while RADIEL and St Carlos, delivering intensive, early, dietitian-led counselling in more selectively enriched populations, achieved significant GDM incidence reductions [21,22,20,25]. This argues against treating "lifestyle intervention" as a single undifferentiated exposure.

Several limitations constrain generalisation. Trials differ in how "standard care" is defined, from no structured advice to fairly detailed printed guidance, so effect size partly reflects how minimal the comparator was [17,5]. Blinding to lifestyle allocation is not feasible, risking performance and detection bias in adherence reporting and insulin-initiation decisions [18]. Prevention-trial populations such as RADIEL, UPBEAT, and LIMIT differ substantially in body mass index, ethnicity, and setting, limiting cross-trial pooling [20,21,22]. The DALI and BAMBINO trials illustrate that even well-designed exercise programmes face adherence challenges, so trial-measured efficacy under supervision may exceed real-world effectiveness at scale [26,24].

Clinically, these findings support structured, dietitian-led, adherence-supported lifestyle intervention as a default component of GDM management, consistent with ACOG and ADA guidance recommending medical nutrition

therapy and structured activity before pharmacological escalation [27,28]. For prevention, evidence favours targeting intensive, early intervention toward women identified as high risk, where RADIEL and St Carlos showed the clearest incidence reductions, rather than uniform low-intensity advice across an unselected population [20,25,6].

## 8. Conclusion

Across randomized trials and systematic reviews spanning both prevention and treatment of gestational diabetes, structured lifestyle intervention combining individualised dietary counselling, prescribed physical activity, and structured monitoring produces a consistent advantage over standard care for maternal glycaemic control, gestational weight gain, and macrosomia or large-for-gestational-age birth, while its effect on caesarean delivery and hypertensive outcomes remains smaller and less consistently reproduced [7,8,17]. The magnitude of benefit appears strongly dependent on programme intensity, adherence support, and timing of initiation rather than on lifestyle counselling in a generic sense, a distinction that should guide how structured programmes are designed and resourced as antenatal services move toward making them a routine rather than exceptional part of GDM care [16,27,28].

## 9. References

1. International Diabetes Federation (2021) IDF Diabetes Atlas, 10th edition. International Diabetes Federation, Brussels.
2. HAPO Study Cooperative Research Group, Metzger, B.E., Lowe, L.P., Dyer, A.R., Trimble, E.R., Chaovarindr, U., et al. (2008) Hyperglycemia and adverse pregnancy outcomes. New England Journal of Medicine, 358(19): 1991-2002. DOI: [10.1056/NEJMoa0707943](https://doi.org/10.1056/NEJMoa0707943)
3. Metzger, B.E., Gabbe, S.G., Persson, B., Buchanan, T.A., Catalano, P.A., Damm, P., et al. (2010) International association of diabetes and pregnancy study groups recommendations on the diagnosis and classification of hyperglycemia in pregnancy. Diabetes Care, 33(3): 676-682. DOI: [10.2337/dc09-1848](https://doi.org/10.2337/dc09-1848)
4. Metzger, B.E., Buchanan, T.A., Coustan, D.R., de Leiva, A., Dunger, D.B., Hadden, D.R., et al. (2007) Summary and recommendations of the Fifth International Workshop-Conference on Gestational Diabetes Mellitus. Diabetes Care, 30(Suppl 2): S251-S260.
5. World Health Organization (2013) Diagnostic Criteria and Classification of Hyperglycaemia First Detected in Pregnancy. World Health Organization, Geneva.
6. Rasmussen, L., Poulsen, C.W., Kampmann, U., Smedegaard, S.B., Ovesen, P.G., Fuglsang, J. (2020) Diet and healthy lifestyle in the management of gestational diabetes mellitus. Nutrients, 12(10): 3050. DOI: [10.3390/nu12103050](https://doi.org/10.3390/nu12103050)
7. Crowther, C.A., Hiller, J.E., Moss, J.R., McPhee, A.J., Jeffries, W.S., Robinson, J.S. (2005) Effect of treatment of gestational diabetes mellitus on pregnancy outcomes. New England Journal of Medicine, 352(24): 2477-2486. DOI: [10.1056/NEJMoa042973](https://doi.org/10.1056/NEJMoa042973)
8. Landon, M.B., Spong, C.Y., Thom, E., Carpenter, M.W., Ramin, S.M., Casey, B., et al. (2009) A multicenter, randomized trial of treatment for mild gestational diabetes. New England Journal of Medicine, 361(14): 1339-1348. DOI: [10.1056/NEJMoa0902430](https://doi.org/10.1056/NEJMoa0902430)
9. Bain, E., Crane, M., Tieu, J., Han, S., Crowther, C.A., Middleton, P. (2015) Diet and exercise interventions for preventing gestational diabetes mellitus. Cochrane Database of Systematic Reviews, (4): CD010443. DOI: [10.1002/14651858.CD010443.pub2](https://doi.org/10.1002/14651858.CD010443.pub2)
10. Han, S., Middleton, P., Crowther, C.A. (2012) Exercise for pregnant women for preventing gestational diabetes mellitus. Cochrane Database of Systematic Reviews, (7): CD009021.
11. Tieu, J., Shepherd, E., Middleton, P., Crowther, C.A. (2017) Dietary advice interventions in pregnancy for preventing

- gestational diabetes mellitus. Cochrane Database of Systematic Reviews, (1): CD006674.
12. Yamamoto, J.M., Kellett, J.E., Balsells, M., Garcia-Patterson, A., Hadar, E., Sola, I., et al. (2018) Gestational diabetes mellitus and diet: a systematic review and meta-analysis of randomized controlled trials examining the impact of modified dietary interventions on maternal glucose control and neonatal birth weight. *Diabetes Care*, 41(7): 1346-1361. DOI: [10.2337/dc18-0102](https://doi.org/10.2337/dc18-0102)
13. Viana, L.V., Gross, J.L., Azevedo, M.J. (2014) Dietary intervention in patients with gestational diabetes mellitus: a systematic review and meta-analysis of randomized clinical trials on maternal and newborn outcomes. *Diabetes Care*, 37(12): 3345-3355. DOI: [10.2337/dc14-1530](https://doi.org/10.2337/dc14-1530)
14. Wang, C., Wei, Y., Zhang, X., Zhang, Y., Xu, Q., Sun, Y., et al. (2017) A randomized clinical trial of exercise during pregnancy to prevent gestational diabetes mellitus and improve pregnancy outcome in overweight and obese pregnant women. *American Journal of Obstetrics and Gynecology*, 216(4): 340-351.
15. Song, C., Li, J., Leng, J., Ma, R.C.W., Yang, X. (2016) Lifestyle intervention can reduce the risk of gestational diabetes: a meta-analysis of randomized controlled trials. *Obesity Reviews*, 17(10): 960-969. DOI: [10.1111/obr.12442](https://doi.org/10.1111/obr.12442)
16. Guo, X.Y., Shu, J., Fu, X.H., Chen, X.P., Zhang, L., Ji, M.X., et al. (2019) Improving the effectiveness of lifestyle interventions for gestational diabetes prevention: a meta-analysis and meta-regression. *BJOG: An International Journal of Obstetrics & Gynaecology*, 126(3): 311-320. DOI: [10.1111/1471-0528.15467](https://doi.org/10.1111/1471-0528.15467)
17. Brown, J., Alwan, N.A., West, J., Brown, S., McKinlay, C.J.D., Farrar, D., Crowther, C.A. (2017) Lifestyle interventions for the treatment of women with gestational diabetes. *Cochrane Database of Systematic Reviews*, (5): CD011970. DOI: [10.1002/14651858.CD011970.pub2](https://doi.org/10.1002/14651858.CD011970.pub2)
18. Hartling, L., Dryden, D.M., Guthrie, A., Muise, M., Vandermeer, B., Donovan, L. (2013) Benefits and harms of treating gestational diabetes mellitus: a systematic review and meta-analysis for the U.S. Preventive Services Task Force and the National Institutes of Health Office of Medical Applications of Research. *Annals of Internal Medicine*, 159(2): 123-129. DOI: [10.7326/0003-4819-159-2-201307160-00661](https://doi.org/10.7326/0003-4819-159-2-201307160-00661)
19. Horvath, K., Koch, K., Jeitler, K., Matyas, E., Bender, R., Bastian, H., et al. (2010) Effects of treatment in women with gestational diabetes mellitus: systematic review and meta-analysis. *BMJ*, 340: c1395. DOI: [10.1136/bmj.c1395](https://doi.org/10.1136/bmj.c1395)
20. Koivusalo, S.B., Rono, K., Klemetti, M.M., Roine, R.P., Lindstrom, J., Erkkola, M., et al. (2016) Gestational diabetes mellitus can be prevented by lifestyle intervention: the Finnish Gestational Diabetes Prevention Study (RADIEL). *Diabetes Care*, 39(1): 24-30. DOI: [10.2337/dc15-0511](https://doi.org/10.2337/dc15-0511)
21. Poston, L., Bell, R., Croker, H., Flynn, A.C., Godfrey, K.M., Goff, L., et al. (2015) Effect of a behavioural intervention in obese pregnant women (the UPBEAT study): a multicentre, randomised controlled trial. *Lancet Diabetes & Endocrinology*, 3(10): 767-777. DOI: [10.1016/S2213-8587\(15\)00227-2](https://doi.org/10.1016/S2213-8587(15)00227-2)
22. Dodd, J.M., Turnbull, D., McPhee, A.J., Deussen, A.R., Grivell, R.M., Yelland, L.N., et al. (2014) Antenatal lifestyle advice for women who are overweight or obese: LIMIT randomised trial. *BMJ*, 348: g1285. DOI: [10.1136/bmj.g1285](https://doi.org/10.1136/bmj.g1285)
23. Luoto, R., Kinnunen, T.I., Aittasalo, M., Kolu, P., Raitanen, J., Ojala, K., et al. (2011) Primary prevention of gestational diabetes mellitus and large-for-gestational-age newborns by lifestyle counseling: a cluster-randomized controlled trial. *PLoS Medicine*, 8(5): e1001036. DOI: [10.1371/journal.pmed.1001036](https://doi.org/10.1371/journal.pmed.1001036)
24. Callaway, L.K., Colditz, P.B., Byrne, N.M., Lingwood, B.E., Rowlands, I.J., Foxcroft, K., et al. (2010) Prevention of gestational diabetes: feasibility issues for an exercise intervention in obese pregnant women. *Diabetes Care*,

33(7): 1457-1459. DOI: [10.2337/dc09-2336](https://doi.org/10.2337/dc09-2336)

25. Assaf-Balut, C., Garcia de la Torre, N., Duran, A., Fuentes, M., Bordiu, E., Del Valle, L., et al. (2017) A Mediterranean diet with additional extra virgin olive oil and pistachios reduces the incidence of gestational diabetes mellitus (GDM): a randomized controlled trial. PLoS ONE, 12(10): e0185873. DOI: [10.1371/journal.pone.0185873](https://doi.org/10.1371/journal.pone.0185873)
26. Simmons, D., Devlieger, R., van Assche, A., Jans, G., Galjaard, S., Corcoy, R., et al. (2017) Association between physical activity and glycemic control in gestational diabetes mellitus: the DALI lifestyle pilot randomized controlled trial. Diabetes Care, 40(4): 469-476.
27. American College of Obstetricians and Gynecologists (2018) ACOG Practice Bulletin No. 190: gestational diabetes mellitus. Obstetrics & Gynecology, 131(2): e49-e64. DOI: [10.1097/AOG.0000000000002501](https://doi.org/10.1097/AOG.0000000000002501)
28. American Diabetes Association (2021) 14. Management of diabetes in pregnancy: standards of medical care in diabetes-2021. Diabetes Care, 44(Suppl 1): S200-S210. DOI: [10.2337/dc21-S014](https://doi.org/10.2337/dc21-S014)
29. Bao, W., Tobias, D.K., Bowers, K., Chavarro, J., Vaag, A., Grunnet, L.G., et al. (2014) Physical activity and sedentary behaviors associated with risk of progression from gestational diabetes mellitus to type 2 diabetes mellitus: a prospective cohort study. JAMA Internal Medicine, 174(7): 1047-1055. DOI: [10.1001/jamainternmed.2014.1795](https://doi.org/10.1001/jamainternmed.2014.1795)
30. Zhang, C., Ning, Y. (2011) Effect of dietary and lifestyle factors on the risk of gestational diabetes: review of epidemiologic evidence. American Journal of Clinical Nutrition, 94(6 Suppl): 1975S-1979S. DOI: [10.3945/ajcn.110.001032](https://doi.org/10.3945/ajcn.110.001032)