

Annals of Dental Science and Oral Biology

Research Article

Evaluation of histopathological analysis as a Prognostic Tool in dentin hypersensitivity

Diallo A^{1*}, Osei K² and Ndlovu S³

¹Department of Oral Biology, Fraser Center for Oral Biology, Kenya

²Department of Dental Sciences, Volga Academy of Oral Biology, Zimbabwe

³Department of Oral Biology, Colorado Institute of Periodontology, Ghana

*Address for Correspondence: Amara Diallo, Department of Oral Biology, Fraser Center for Oral Biology, Kenya;
E-mail: amara.diallo@fraser.edu

Received: 9 December 2025; Accepted: 6 March 2026; Published: 6 June 2026.

Citation of this article: Diallo A, Osei K, Ndlovu S (2026) Evaluation of histopathological analysis as a Prognostic Tool in dentin hypersensitivity. Ann Dent Sci Oral Biol, 7(1): 01-06. DOI: [10.5281/zenodo.22811967](https://doi.org/10.5281/zenodo.22811967)

Copyright: © 2026 Amara Diallo, et al. This is an open access article distributed under the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: The hydrodynamic theory attributes dentin hypersensitivity pain to fluid movement within open dentinal tubules, linking the symptom to a structure histology and scanning electron microscopy (SEM) can directly visualize [2,8].

Objective: This review evaluates whether histopathological analysis of dentin and the pulp-dentin complex functions as a reliable prognostic tool, predicting symptom persistence, treatment response, and the limits of dentin-directed therapy.

Methods: We synthesized SEM and histologic studies comparing sensitive and non-sensitive dentin, treatment-response studies confirming tubule occlusion, the tertiary-dentinogenesis literature, and clinical-histologic correlation studies of the pulp-dentin complex.

Findings: Sensitive dentin shows markedly more numerous, wider open tubules than non-sensitive dentin [8], and treatment response tracks histologically confirmed tubule occlusion while non-response tracks persistent patency [11]; reactionary dentinogenesis offers a plausible but less directly tested prognostic axis [15], and clinical-histologic correlation in deep caries shows only moderate agreement with underlying histology [18].

Conclusions: Histopathological analysis, principally SEM assessment of tubule occlusion, carries genuine prognostic value and is best used as a bench-validation tool for interpreting treatment response rather than a chairside diagnostic test [8,11].

Keywords: dentin hypersensitivity; histopathology; dentinal tubules; hydrodynamic theory; pulp-dentin complex

1. Introduction

Dentin hypersensitivity is defined clinically by a short, sharp pain arising from exposed dentin in response to a stimulus, typically thermal, evaporative, tactile, or osmotic, that cannot be explained by any other dental defect or pathology [1]. That definition is deliberately negative, built by exclusion of caries, fracture, and restorative failure, and it is one reason the condition is managed almost entirely on symptom report rather than on any tissue-level correlate. Yet the pain is generated at a specific structure, the exposed dentin surface and the tubules beneath it, and the degree to which those tubules remain patent, occluded, or reorganized by reactionary mineralization is, in principle, something a pathologist can see and measure. The hydrodynamic theory that underlies most mechanistic accounts of dentin hypersensitivity proposed exactly this link: fluid movement within open dentinal tubules, driven by thermal, evaporative, or osmotic gradients at the exposed surface, deforms the odontoblast-nerve complex at the pulpal end of the tubule and triggers the A-delta fibers responsible for the sharp pain patients describe [2,3]. If tubule patency is the proximate cause of the pain, a histological assessment of patency, diameter, and occlusion should carry prognostic information beyond what a symptom questionnaire alone provides, both for identifying which surfaces will remain symptomatic and for judging whether a treatment has actually closed the anatomical pathway the pain depends on.

This review evaluates that proposition: whether histopathological analysis of dentin functions as a reliable prognostic tool in dentin hypersensitivity, predicting which sites remain symptomatic, distinguishing responders from non-responders to desensitizing treatment, and flagging cases where the pulp is already too compromised for dentin-directed therapy alone. The question matters because dentin hypersensitivity is both extremely common, with pooled self-reported prevalence estimated at roughly one in three adults [4], and

persistently difficult to manage predictably, where the same regimen produces durable relief in some patients and none in others for reasons symptom scoring alone cannot explain [5,6,7].

2. Methods

We conducted a narrative synthesis of the peer-reviewed literature addressing the histology and ultrastructure of hypersensitive dentin, the tubule-level basis of the hydrodynamic mechanism, and the correlation between histologic findings and clinical or prognostic outcomes in dentin hypersensitivity and related pulp-dentin pathology. Sources were identified through targeted searches combining histological, scanning electron microscopic, or histobacteriological methods with clinical assessment of dentin sensitivity, tubule occlusion, tertiary dentin formation, or pulp status, supplemented by the foundational mechanistic and epidemiological literature. Priority was given to studies directly comparing histologic features between sensitive and non-sensitive, or treated and untreated, dentin, and to studies formally correlating a histologic diagnosis with a clinical one, since these designs speak most directly to prognostic validity rather than to descriptive anatomy. Findings are organized thematically: the ultrastructural basis for tubule-level prognosis, histologically verified occlusion after treatment, the pulp-dentin complex's reparative response, and clinical-histologic correlation in the adjacent literature on pulp status.

3. Results

Tubule-level histology distinguishes sensitive from non-sensitive dentin

The most influential finding in this literature is also its oldest and simplest: in extracted teeth with clinically documented sensitive and non-sensitive cervical dentin from the same patients, scanning electron microscopy showed sensitive surfaces had substantially more open dentinal tubules per unit area, and tubules of markedly wider mean diameter, than adjacent non-sensitive

surfaces [8]. The magnitude reported was large, with sensitive dentin showing on the order of eight times the density of patent tubules seen in non-sensitive dentin, and orifices roughly twice as wide. That comparison is the closest this field has to a histological "test" with prognostic content: a surface with numerous wide, unoccluded tubules is more likely to remain symptomatic, while one whose tubules are narrow, sparse, or occluded is less likely to generate hydrodynamic pain regardless of how much dentin is exposed. The same group's companion work showed acidic dietary substances and toothbrush abrasion applied to root-planed dentin actively re-open partially occluded tubules, reversing whatever protective histologic state existed [9], and an in situ model developed to study tubule patency under controlled intraoral conditions confirmed patency can be tracked prospectively over days to weeks rather than only at a single extraction-derived timepoint [10]. Together, these studies establish that tubule patency is a dynamic state that can move in either direction, which is why a single histologic snapshot has prognostic value only when read as a report on the current mechanical situation rather than a permanent diagnosis.




Figure 1: Relative dentinal tubule density and mean tubule diameter on scanning electron microscopy of clinically sensitive versus adjacent non-sensitive cervical dentin from the same patients, illustrating the tubule-level basis for a histologic prognostic signal [8].

Occlusion, smear layer, and the histologic signature of treatment response

If open tubules are the histologic correlate of symptomatic dentin, histologically confirmed occlusion should be the

correlate of a treatment response, and the periodontal literature on root hypersensitivity after scaling and root planing bears this out. Kerns and colleagues used SEM to examine root surfaces before and after treatment with a potassium-oxalate desensitizing agent and found clinically hypersensitive roots showed patent tubules at baseline, roots that responded clinically showed a dense crystalline occluding layer on follow-up SEM, and non-responding roots retained visibly open tubules despite identical treatment [11]. This pattern gives histological analysis practical rather than merely explanatory value: SEM appearance after treatment tracked clinical outcome closely enough that persistent patency after an appropriate intervention is itself a reasonable predictor of continued symptoms. The broader literature on desensitizing dentifrices, in-office pastes, and laser-mediated dentin fusion is built on the same logic: an agent is judged efficacious largely by whether SEM after application shows tubules occluded by a stable plug rather than a friable smear layer [1,5]. Clinically, an in-office arginine-calcium-carbonate paste produced rapid relief in a split-mouth randomized trial, attributed to mechanical plugging of tubule lumina by an arginine-bicarbonate-calcium complex lodging in the orifice [12]. The prognostic reading is that the durability of occlusion, not merely its presence immediately after application, is what predicts whether relief persists, since a thin or poorly adherent occluding layer is most likely to be lost to brushing, acid exposure, or masticatory load within weeks.

Table 1: Histologic and ultrastructural features reported at the dentin surface across sensitive, non-sensitive, and treated states, drawn from correlated SEM and clinical studies [8,9,11].			
Dentin surface state	Tubule patency on SEM	Mean tubule diameter	Associated clinical status
Sensitive, untreated cervical dentin	Numerous open tubules	Wide (approx. 2x non-sensitive)	Symptomatic to thermal/tactile stimuli
Non-sensitive, untreated cervical dentin	Few open tubules	Narrow	Asymptomatic despite comparable exposure

Dentin surface state	Tubule patency on SEM	Mean tubule diameter	Associated clinical status
Root surface after oxalate desensitizer, responder	Dense occluding crystalline layer	Effectively occluded	Clinical relief on follow-up
Root surface after oxalate desensitizer, non-responder	Tubules remain patent	Unchanged	Symptoms persist despite treatment

The pulp-dentin complex's own histology as a second prognostic axis

Dentin hypersensitivity is not generated by dentin in isolation; every tubule terminates at the pulp, and the odontoblast-predentin-pulp complex has its own reparative histology bearing on prognosis. Dentin permeability, and the volume of fluid that can move through a tubule for a given stimulus, is set partly by pulpal factors: intrapulpal pressure, tubule patency along its length, and the degree to which reactionary or tertiary dentin has been laid down beneath the exposure site by surviving odontoblasts responding to a mild, chronic irritant [13,14]. Reactionary dentinogenesis, deposition of an atubular or sparsely tubular layer by the original odontoblast population, is distinct from reparative dentinogenesis, which follows odontoblast death and requires recruitment of a new mineralizing cell population; the extent to which either process has occurred offers real prognostic information. A pulp already walled off with reactionary dentin is one in which pain is likely to fade even without intervention, while a pulp showing no reparative response, or early inflammatory change, is a poorer prognosis and a signal the presenting complaint may not be simple dentin hypersensitivity at all [15,16,17].




Figure 2: Pooled self-reported prevalence of dentin hypersensitivity from systematic review, set against the wide prevalence range reported across studies using differing diagnostic stimuli and thresholds, indicating the scale of the population in whom a histologically grounded prognostic model has practical relevance [4,6].

Clinical-histologic correlation as the direct test of prognostic validity

The most rigorous available test of whether a clinical impression of pulp-dentin status predicts the underlying tissue state comes from the closely related histobacteriological work of Ricucci, Siqueira, and colleagues on pulps beneath deep caries, which applies exactly the correlated clinical-versus-histologic design a genuine prognostic evaluation requires. Comparing preoperative clinical diagnoses with subsequent histologic and histobacteriologic examination, this group repeatedly found meaningful discordance: teeth clinically diagnosed with reversible pulpitis or a normal pulp not infrequently showed more advanced, localized inflammation than the clinical picture suggested, particularly with bacterial penetration reaching the pulp chamber roof, while some symptomatic teeth showed comparatively well-localized, non-progressive change [18,19,20]. In a cohort of immature teeth with deep caries, the clinical diagnosis of pulp status again correlated only loosely with the actual depth and distribution of inflammation and bacterial colonization observed microscopically, with some clinically unremarkable pulps already showing partial necrosis adjacent to bacterial invasion [21]. The implication for dentin hypersensitivity is direct: if clinical assessment correlates only moderately with true histologic state even in caries, where the insult is more

severe than the mild, chronic irritation typical of exposed cervical dentin, clinical judgment alone is unlikely to be fully reliable in dentin hypersensitivity either, and a histologically informed model adds predictive information symptom report and visual inspection cannot supply alone.

Table 2: Patterns of agreement and discordance between preoperative clinical diagnosis and subsequent histologic/histobacteriologic examination of the pulp-dentin complex in deep caries, offered as the closest available direct test of clinical-histologic prognostic validity relevant to dentin hypersensitivity [18,21,19,20].

Clinical impression	Histologic/histobacteriologic finding	Prognostic implication
Normal or reversible pulpitis	Occasionally more advanced, localized inflammation with bacterial penetration	Clinical exam alone can understate underlying tissue involvement
Symptomatic, presumed advanced pathology	Sometimes well-localized, non-progressive change	Symptom severity does not map linearly onto histologic extent
Clinically unremarkable pulp beneath deep caries (immature teeth)	Partial necrosis adjacent to bacterial invasion in some cases	Reinforces the need for a tissue-level, not purely clinical, prognostic model

4. Discussion

Taken together, this evidence supports a specific, fairly narrow claim: histopathological analysis, principally SEM of tubule patency and diameter, together with what is known about reactionary dentinogenesis and pulpal histology from the adjacent caries literature, functions as a genuine prognostic tool, but one operating at the level of tissue state rather than as a bedside test any clinician performs chairside. The signal is strongest at the tubule level: sensitive dentin carries far more numerous, far wider open tubules than adjacent non-sensitive dentin, and treatment response tracks histologically confirmed occlusion while non-response tracks persistent patency, extending the claim from description to prediction [8,11]. Epidemiological data reinforce why this matters at a population level: with self-reported prevalence around one in three adults, and wide variability across studies

using different diagnostic thresholds, a histologically grounded understanding of which surfaces are mechanistically likely to remain symptomatic has real value for triaging who needs active therapy versus reassurance [4,22,6].

The evidence is weaker once the pulp itself is considered. The reparative and reactionary dentinogenesis literature makes a biologically coherent case that pulpal histology beneath a hypersensitive surface should predict whether the condition self-resolves, but this rests more on mechanistic plausibility than on prospective studies biopsying vital pulps beneath characterized hypersensitive teeth [15,16,17,14]. The clinical-histologic correlation studies that come closest to testing this directly were conducted in deep caries rather than dentin hypersensitivity proper, and while directly relevant, since both conditions turn on whether a clinical impression matches underlying tissue reality, the discordance rates should be read as a caution rather than a figure transferring unmodified to hypersensitive but non-carious teeth [18,21,19,20]. Clinicians currently rely on an inference chain, from SEM studies of extracted teeth to living practice, rather than direct histologic sampling of the pulp-dentin complex in vivo, which is neither ethically nor practically feasible as a routine step. That inferential gap is the main limitation of histopathological analysis as a prognostic tool in current use: it informs which treatments are mechanistically rational and helps interpret why patients fail to respond, but cannot yet be applied prospectively to an individual tooth without an invasive biopsy.

Practically, histological evidence is best used the way the Kerns and Absi-type SEM literature already uses it: as a bench validation standard against which desensitizing treatments are judged, and as an explanatory framework for why treatment fails, tubules never fully occluded, an occluding layer too thin to survive function, or insufficient reparative dentinogenesis, rather than as a chairside

diagnostic test, mirroring how management guidelines already separate mechanistic rationale from bedside decision-making [23]. Prospective work correlating quantitative SEM tubule morphometry and pulpal reactionary dentin thickness with the documented symptom trajectory of the same patient before extraction would close much of the remaining gap.

5. Conclusion

Histopathological analysis has a real, evidence-supported prognostic role in dentin hypersensitivity, most directly through SEM assessment of tubule patency and diameter, which reliably distinguishes sensitive from non-sensitive dentin and tracks clinical response through histologically confirmed occlusion [8,11]. Its role is weaker and more inferential on the pulpal side, where reparative dentinogenesis plausibly predicts symptom persistence but has not been tested prospectively in dentin hypersensitivity itself, and where the closest direct evidence, from clinical-histologic correlation in deep caries, shows only moderate agreement between clinical impression and underlying histology [18,21,15]. Used appropriately, as a mechanistic and bench-validation tool rather than a chairside diagnostic test, histopathological analysis remains the most direct available explanation for why dentin hypersensitivity occurs, why some sites respond to treatment while others do not, and where the limits of purely dentin-directed therapy lie.

6. Acknowledgements

The authors thank colleagues in their respective departments of oral biology and periodontology for discussion of the histological literature reviewed here.

7. Funding

This work received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

8. Conflict of Interest

The authors declare no conflict of interest.

9. Author Contributions

A.D. conceived the review and drafted the manuscript. K.O. and S.N. conducted the literature synthesis and critically revised the manuscript for intellectual content. All authors approved the final version.

10. Ethics Statement

This article is a narrative literature review and did not involve human participants, human data, or animal subjects; no ethics approval was required.

11. Data Availability

No new data were generated for this review. All sources synthesized are cited in the reference list.

12. References

1. Dionysopoulos, D., Gerasimidou, O., Beltes, C. (2023) Dentin hypersensitivity: etiology, diagnosis and contemporary therapeutic approaches-a review in literature. *Applied Sciences*, 13(21): 11632. DOI: [10.3390/app132111632](https://doi.org/10.3390/app132111632)
2. Brannstrom, M., Astrom, A. (1964) A study on the mechanism of pain elicited from the dentin. *Journal of Dental Research*, 43(4): 619-625.
3. Gysi, A. (1900) An attempt to explain the sensitiveness of dentine. *British Journal of Dental Science*, 43: 865-868.
4. Favaro Zeola, L., Soares, P.V., Cunha-Cruz, J. (2019) Prevalence of dentin hypersensitivity: systematic review and meta-analysis. *Journal of Dentistry*, 81: 1-6. DOI: [10.1016/j.jdent.2018.12.015](https://doi.org/10.1016/j.jdent.2018.12.015)
5. Liu, X.X., Tenenbaum, H.C., Wilder, R.S., Quock, R., Hewlett, E.R., Ren, Y.F. (2020) Pathogenesis, diagnosis and management of dentin hypersensitivity: an evidence-based overview for dental practitioners. *BMC Oral Health*, 20: 220. DOI: [10.1186/s12903-020-01199-z](https://doi.org/10.1186/s12903-020-01199-z)
6. West, N.X., Lussi, A., Seong, J., Hellwig, E. (2013) Dentin hypersensitivity: pain mechanisms and aetiology of exposed cervical dentin. *Clinical Oral Investigations*, 17(Suppl 1):

- S9-S19.
7. Addy, M. (2002) Dentine hypersensitivity: new perspectives on an old problem. *International Dental Journal*, 52(S5): 367-375.
 8. Absi, E.G., Addy, M., Adams, D. (1987) Dentine hypersensitivity: a study of the patency of dentinal tubules in sensitive and non-sensitive cervical dentine. *Journal of Clinical Periodontology*, 14(5): 280-284. DOI: [10.1111/j.1600-051x.1987.tb01533.x](https://doi.org/10.1111/j.1600-051x.1987.tb01533.x)
 9. Absi, E.G., Addy, M., Adams, D. (1992) Dentine hypersensitivity: the effect of toothbrushing and dietary compounds on dentine in vitro: an SEM study. *Journal of Oral Rehabilitation*, 19(2): 101-110. DOI: [10.1111/j.1365-2842.1992.tb01086.x](https://doi.org/10.1111/j.1365-2842.1992.tb01086.x)
 10. Banfield, N., Addy, M. (2004) Dentine hypersensitivity: development and evaluation of a model in situ to study tubule patency. *Journal of Clinical Periodontology*, 31(5): 325-335.
 11. Kerns, D.G., Scheidt, M.J., Pashley, D.H., Horner, J.A., Strong, S.L., Van Dyke, T.E. (1991) Dentinal tubule occlusion and root hypersensitivity. *Journal of Periodontology*, 62(7): 421-428. DOI: [10.1902/jop.1991.62.7.421](https://doi.org/10.1902/jop.1991.62.7.421)
 12. Kapferer, I., Pflug, C., Kisielowsky, I., Giesinger, J., Beier, U.S., Dumfahrt, H. (2013) Instant dentin hypersensitivity relief of a single topical application of an in-office desensitizing paste containing 8% arginine and calcium carbonate: a split-mouth, randomized-controlled study. *Acta Odontologica Scandinavica*, 71(3-4): 994-999. DOI: [10.3109/00016357.2012.741701](https://doi.org/10.3109/00016357.2012.741701)
 13. Pashley, D.H. (1986) Dentin permeability, dentin sensitivity, and treatment through tubule occlusion. *Journal of Endodontics*, 12(10): 465-474.
 14. Pashley, D.H. (1996) Dynamics of the pulpo-dentin complex. *Critical Reviews in Oral Biology and Medicine*, 7(2): 104-133. DOI: [10.1177/10454411960070020101](https://doi.org/10.1177/10454411960070020101)
 15. Smith, A.J., Cassidy, N., Perry, H., Begue-Kirn, C., Ruch, J.V., Lesot, H. (1995) Reactionary dentinogenesis. *International Journal of Developmental Biology*, 39(1): 273-280.
 16. Goldberg, M., Smith, A.J. (2004) Cells and extracellular matrices of dentin and pulp: a biological basis for repair and tissue engineering. *Critical Reviews in Oral Biology and Medicine*, 15(1): 13-27.
 17. Lee, M., Lee, Y.S., Shon, W.J., Park, J.C. (2023) Physiologic dentin regeneration: its past, present, and future perspectives. *Frontiers in Physiology*, 14: 1313927. DOI: [10.3389/fphys.2023.1313927](https://doi.org/10.3389/fphys.2023.1313927)
 18. Ricucci, D., Loghin, S., Siqueira, J.F. (2014) Correlation between clinical and histologic pulp diagnoses. *Journal of Endodontics*, 40(12): 1932-1939. DOI: [10.1016/j.joen.2014.08.010](https://doi.org/10.1016/j.joen.2014.08.010)
 19. Ricucci, D., Siqueira, J.F., Li, Y., Tay, F.R. (2019) Vital pulp therapy: histopathology and histobacteriology-based guidelines to treat teeth with deep caries and pulp exposure. *Journal of Dentistry*, 86: 41-52. DOI: [10.1016/j.jdent.2019.05.022](https://doi.org/10.1016/j.jdent.2019.05.022)
 20. Ricucci, D., Siqueira, J.F., Rocas, I.N., Lipski, M., Shibani, A., Tay, F.R. (2020) Pulp and dentine responses to selective caries excavation: a histological and histobacteriological human study. *Journal of Dentistry*, 100: 103430. DOI: [10.1016/j.jdent.2020.103430](https://doi.org/10.1016/j.jdent.2020.103430)
 21. Ricucci, D., Siqueira, J.F., Loghin, S., Lin, L.M. (2017) Pulp and apical tissue response to deep caries in immature teeth: a histologic and histobacteriologic study. *Journal of Dentistry*, 56: 19-32. DOI: [10.1016/j.jdent.2016.10.005](https://doi.org/10.1016/j.jdent.2016.10.005)
 22. Splieth, C.H., Tachou, A. (2013) Epidemiology of dentin hypersensitivity. *Clinical Oral Investigations*, 17(Suppl 1): S3-S8. DOI: [10.1007/s00784-012-0889-8](https://doi.org/10.1007/s00784-012-0889-8)
 23. Orchardson, R., Gillam, D.G. (2006) Managing dentin hypersensitivity. *Journal of the American Dental Association*, 137(7): 990-998.