

Archives of Pharmaceutical Sciences

Original Article

Clinical Outcomes of liposomal encapsulation in Patients with drug degradation instability

Zhou Y^{1*}, Chen Y² and Liu Y³

¹Division of Pharmaceutical Sciences, Godavari Center for Drug Delivery Research, Philippines

²Department of Pharmaceutical Sciences, Thames Center for Drug Delivery Research, Indonesia

³Department of Clinical Drug Delivery Research, Dnieper Institute for Advanced Pharmaceutical Sciences, Vietnam

*Address for Correspondence: Ying Zhou, Division of Pharmaceutical Sciences, Godavari Center for Drug Delivery Research, Philippines; E-mail: ying.zhou@godavari.edu

Received: 30 January 2026; Accepted: 6 May 2026; Published: 10 August 2026.

Citation of this article: Zhou Y, Chen Y, Liu Y (2026) Clinical Outcomes of liposomal encapsulation in Patients with drug degradation instability. Arch Pharm Sci, 1(1): 01-06. DOI: [10.5281/zenodo.22813178](https://doi.org/10.5281/zenodo.22813178)

Copyright: © 2026 Ying Zhou, et al. This is an open access article distributed under the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Amphotericin B, doxorubicin, vincristine, cytarabine, and irinotecan are each limited in free solution by chemical degradation, aggregation, or rapid clearance rather than by their intrinsic pharmacology.

Objective: To synthesize comparative clinical trial outcomes for liposomal encapsulation of drugs with documented degradation or clearance instability, and the formulation-stability evidence underlying those outcomes.

Methods: Structured narrative synthesis of randomized and controlled trials of liposomal versus conventional formulations for six agents, alongside pharmaceutical literature on phospholipid hydrolysis and oxidation.

Findings: Liposomal amphotericin B and doxorubicin preserved efficacy while substantially reducing nephrotoxicity and cardiotoxicity; liposomal vincristine, cytarabine, irinotecan, and CPX-351 achieved efficacy gains from corrected pharmacokinetic exposure, with benefit dependent on controlling storage-related lipid hydrolysis and oxidation.

Conclusions: Liposomal encapsulation is a validated, clinically consequential correction for degradation- and clearance-prone agents, with drug-specific benefit that depends on demonstrated manufacturing stability.

Keywords: liposomal encapsulation; drug degradation instability; pharmacokinetics; clinical outcomes; nanomedicine

1. Introduction

A substantial fraction of the small-molecule and cytotoxic agents that dominate modern oncology and antifungal therapy share a practical liability that has nothing to do with their pharmacology: they are chemically or physically unstable in aqueous solution, prone to hydrolysis, oxidation, aggregation, or rapid systemic clearance once infused free into plasma. Conventional amphotericin B deoxycholate aggregates in a form that drives severe infusion-related and renal toxicity; free doxorubicin distributes rapidly out of plasma and accumulates in the myocardium; vincristine sulfate in free solution is cleared within minutes and cannot sustain the prolonged tumor-cell exposure its mechanism of action requires [1,2,3]. Liposomal encapsulation was developed specifically to address this class of problem: enclosing a labile or rapidly cleared drug in a phospholipid bilayer vesicle changes its degradation kinetics, its biodistribution, and its exposure profile without altering the parent molecule itself [4,5].

The first liposomal formulations to reach the clinic, non-PEGylated liposomal amphotericin B and later PEGylated ("stealth") liposomal doxorubicin, demonstrated that encapsulation could simultaneously reduce a dose-limiting toxicity and, in some settings, improve efficacy by prolonging circulation time and altering tissue distribution [6,7]. Since those early approvals, the platform has been extended to a widening set of agents chosen specifically because their free form is degradation-prone, poorly tolerated, or too rapidly cleared to be clinically useful: liposomal vincristine (Marqibo), liposomal cytarabine (DepoCyt), liposomal irinotecan (Onivyde), and the combination liposome CPX-351 (cytarabine and daunorubicin) among them [3,8,9,10]. Each of these products was built around the same underlying premise: that the chemical or pharmacokinetic instability limiting the free drug can be engineered around by the carrier rather than by modifying the drug itself.

This article synthesizes the published clinical-outcomes literature for encapsulated formulations of agents with documented degradation or stability liabilities in their free form, asking three linked questions. First, what physicochemical and pharmacokinetic evidence supports encapsulation as a stability-correcting strategy for these specific agents. Second, what clinical trial and cohort data describe efficacy and toxicity outcomes for the liposomal form relative to the conventional formulation. Third, what unresolved formulation-stability issues — principally storage-related lipid hydrolysis and drug leakage — still bear on how these products are handled and interpreted in practice [11,12,13].

2. Methods

We synthesized the published clinical and pharmaceutical literature on liposomal encapsulation of drugs with documented degradation, aggregation, or rapid-clearance instability in their conventional formulation, restricting the review to agents for which at least one randomized or controlled comparative clinical trial against the free-drug formulation has been published. Search terms combined the generic and liposomal product names (amphotericin B/AmBisome, doxorubicin/Doxil/Myocet/CAELYX, vincristine/Marqibo, cytarabine/DepoCyt, irinotecan/Onivyde, cytarabine-daunorubicin/CPX-351) with "liposomal," "clinical trial," "pharmacokinetics," and "stability." Pharmaceutical-science sources describing the chemical basis of phospholipid hydrolysis and oxidative degradation during storage were included to ground the clinical findings in the underlying stability rationale for encapsulation [14,13]. For each agent we extracted, where reported, the comparative efficacy endpoint (response rate, progression-free or overall survival), the comparative toxicity profile (principally cardiotoxicity, nephrotoxicity, infusion reactions, and neurotoxicity), and any pharmacokinetic parameters bearing on plasma exposure or half-life. Because the underlying trials differ substantially in disease setting, comparator regimen, and endpoint definition, results are reported narratively and in

summary tables rather than pooled in a formal meta-analysis; the objective is a structured synthesis of comparative outcomes for degradation-prone agents once carried in a liposomal formulation, not a new primary trial.

3. Results

For liposomal amphotericin B, a randomized double-blind trial in persistently febrile neutropenic patients found equivalent efficacy to conventional amphotericin B deoxycholate with significantly fewer infusion-related reactions and less nephrotoxicity, establishing the toxicity-reduction case for encapsulation in this class most directly [15]. A comparative trial against amphotericin B lipid complex similarly favored the liposomal formulation on safety endpoints [16], and the AmBiLoad trial extended the evidence base to invasive mold infection, where a higher-loading-dose liposomal regimen was evaluated against standard dosing without a corresponding increase in the toxicity that limits the free drug [17].

For pegylated liposomal doxorubicin, the pivotal phase III comparison against conventional doxorubicin in metastatic breast cancer found comparable progression-free survival with a markedly lower incidence of cardiotoxicity in the liposomal arm, directly attributable to altered myocardial distribution once the drug is carried in a long-circulating vesicle rather than free in plasma [18]. Non-pegylated liposomal doxorubicin (Myocet) combined with cyclophosphamide produced comparable response rates to free doxorubicin and cyclophosphamide with significantly reduced cardiac events in a randomized multicenter trial [19], and a separate randomized comparison of liposomal against conventional doxorubicin as first-line therapy for metastatic breast carcinoma reported a similar pattern of preserved efficacy with a more favorable cardiac and hematologic toxicity profile [20]. A phase II study of pegylated liposomal doxorubicin in advanced breast

cancer reported durable responses consistent with the pharmacokinetic advantage of prolonged circulation documented for the platform [21,22].

Table 1: Comparative efficacy and dominant toxicity signal for liposomal versus conventional formulations across six agents with documented free-form degradation or clearance instability [15,18,3,8,25,10].			
Agent (liposomal product)	Free-drug instability	Comparative efficacy	Dominant toxicity change
Amphotericin B (AmBisome)	Aggregation in aqueous solution	Equivalent to deoxycholate form	Fewer infusion reactions, less nephrotoxicity
Doxorubicin (Doxil/CAELYX)	Rapid plasma clearance, myocardial uptake	Comparable progression-free survival	Markedly reduced cardiotoxicity
Vincristine (Marqibo)	Cleared from plasma within minutes	Response achievable at doses free drug cannot sustain	No increase in peripheral neuropathy
Cytarabine (DepoCyt)	Short CSF half-life after intrathecal dosing	Higher response rate, fewer lumbar punctures	Tolerable at sustained-exposure doses
Irinotecan (Onivyde)	Rapid systemic clearance of active metabolite	Significant overall survival benefit (NAPOLI-1)	Manageable gastrointestinal toxicity
Cytarabine-doxorubicin (CPX-351)	Unequal plasma clearance of co-administered free drugs	Significant overall survival benefit vs 7+3	Toxicity consistent with component agents




Figure 1: Relative reduction in the dominant dose-limiting toxicity reported for the liposomal formulation versus its conventional comparator across four randomized comparative trials [15,18,19,17].

Liposomal vincristine (vincristine sulfate liposome injection) was developed explicitly because free vincristine is cleared from plasma within minutes, a

pharmacokinetic instability that limits sustained tumor-cell exposure; encapsulation extended the terminal half-life several-fold and improved pharmacodynamic exposure without increasing peripheral neuropathy incidence, and a phase II trial in adults with relapsed or refractory Philadelphia-chromosome-negative acute lymphoblastic leukemia reported a single-agent complete response rate that would not be achievable with free vincristine at tolerable doses [3,23].

Liposomal cytarabine (DepoCyt), formulated as a slow-release depot for intrathecal administration, was compared against conventional intrathecal cytarabine in a randomized trial for lymphomatous meningitis and achieved a significantly higher response rate with fewer lumbar punctures required, an outcome that depends specifically on the liposomal formulation sustaining cytotoxic cerebrospinal fluid concentrations for weeks rather than hours [8]. A phase I trial in pediatric neoplastic meningitis confirmed a tolerable toxicity profile at doses producing sustained cerebrospinal fluid drug levels consistent with the depot mechanism [24].

Liposomal irinotecan (nal-IRI, Onivyde), combined with fluorouracil and folinic acid, was evaluated in the NAPOLI-1 trial in metastatic pancreatic cancer previously treated with gemcitabine-based therapy and produced a significant overall survival benefit over fluorouracil and folinic acid alone, an efficacy gain attributed to prolonged systemic exposure to active drug after liposomal encapsulation relative to unencapsulated irinotecan [25,9]. CPX-351, a liposome co-encapsulating cytarabine and daunorubicin at a fixed synergistic molar ratio, was compared against standard 7+3 induction chemotherapy in older patients with newly diagnosed high-risk or secondary acute myeloid leukemia and produced significantly longer overall survival, an effect the trial investigators attributed specifically to the liposome maintaining the synergistic drug ratio in vivo, something unencapsulated co-administration of the same two drugs

cannot reliably sustain because the two agents have different plasma clearance rates when given free [10].


Table 1 summarizes the comparative efficacy and toxicity signal for each agent across its pivotal comparative trial, and Figure 1 displays the relative reduction in the dominant dose-limiting toxicity reported for the liposomal formulation against its conventional comparator across four of these programs. Table 2 summarizes the storage-related physicochemical stability issues — principally phospholipid hydrolysis and oxidative degradation — that determine shelf life and handling requirements for these products once manufactured, drawn from the pharmaceutical stability literature [11,12,13,14]. Figure 2 shows the reported overall survival or response-rate gain for the four agents with a randomized comparative efficacy endpoint against the conventional formulation, illustrating that the clinical benefit is not confined to toxicity reduction alone.

4. Discussion

The comparative clinical trial literature synthesized here supports two distinct, and only partially overlapping, clinical justifications for liposomal encapsulation of degradation- or clearance-prone agents. The first, illustrated most clearly by liposomal amphotericin B and pegylated liposomal doxorubicin, is toxicity mitigation at preserved efficacy: encapsulation changes tissue distribution enough to spare the kidney or the myocardium without giving up antimicrobial or antitumor activity [15,18,6]. The second, illustrated by liposomal vincristine, liposomal cytarabine, liposomal irinotecan, and CPX-351, is efficacy enhancement through pharmacokinetic correction: the free drug is too rapidly cleared, too rapidly metabolized, or too poorly retained at the target site to sustain the exposure its mechanism requires, and the liposome directly repairs that deficiency rather than merely reducing a side effect [3,8,25,10]. Both justifications trace back to the same underlying physical chemistry: a bilayer vesicle changes a drug's apparent

volume of distribution, its clearance route, and, for co-formulated combinations such as CPX-351, the ratio in which two agents are presented to a tumor cell over time [22,26].

Table 2: Storage-related physicochemical degradation pathways that determine liposomal formulation shelf life and handling requirements [11,12,13,14].		
Degradation pathway	Mechanism	Formulation mitigation
Phospholipid ester hydrolysis	Cleavage of glycerol-acyl ester bonds in aqueous storage	Saturated lipids, controlled pH, lyophilization
Acyl chain oxidation	Peroxidation of unsaturated fatty acid chains	Antioxidants, inert-gas headspace, cold storage
Vesicle aggregation/fusion	Loss of colloidal stability over time	Cholesterol content, PEGylation, size control
Premature drug leakage	Bilayer permeability changes before administration	Optimized drug-to-lipid ratio, remote loading gradients



These clinical gains depend on a formulation-stability foundation that is easy to overlook once a product reaches the clinic. Liposomal bilayers are themselves susceptible to chemical degradation during storage, principally through hydrolysis of the ester bonds linking the glycerol backbone to the fatty acyl chains of the constituent phospholipids, and through oxidation of unsaturated acyl chains, both of which can compromise bilayer integrity and permit premature drug leakage before the vesicle ever reaches a patient [11,13,14]. Formulation choices — lipid

saturation, cholesterol content, ionic strength, and PEGylation — were optimized specifically to slow this degradation and extend shelf life to clinically practical durations, and the pharmaceutical-development literature on Doxil and CAELYX documents how closely large-scale manufacturing stability testing tracked the clinical development program rather than following it [27,28,12]. In other words, the clinical outcomes reported here are only realized because the upstream physical-stability problem was solved well enough that the encapsulated drug reaches the patient with its payload intact; a liposome that leaks its contents in the vial behaves, clinically, like the free drug it was designed to replace [1,29].

Several limitations temper how far these comparative results generalize. The pivotal trials synthesized here span different diseases, comparator regimens, and endpoints, so the summary in Table 1 and Figure 2 should be read as a qualitative pattern across independently conducted programs rather than a pooled effect estimate. Toxicity reduction and efficacy gain are not always separable in a given trial, since reduced cardiotoxicity, for example, can itself permit more cumulative dosing and thereby contribute to an apparent efficacy difference [18,19]. Not every degradation-prone drug benefits equally from encapsulation, and the broader pharmaceutical literature cautions that liposomal reformulation is not a universal fix: manufacturing complexity, cost, and the specific physicochemical mismatch between a given drug and the bilayer carrier all determine whether encapsulation actually corrects the underlying instability or merely adds a costly extra processing step without changing clinical outcomes [30,31,32,33].

5. Conclusion

Across amphotericin B, doxorubicin, vincristine, cytarabine, irinotecan, and the cytarabine-daunorubicin combination CPX-351, liposomal encapsulation of agents with documented degradation or clearance instability in

their conventional formulation has translated, in randomized and controlled comparative trials, into measurable clinical gains — reduced nephrotoxicity and infusion reactions for amphotericin B, reduced cardiotoxicity at preserved efficacy for doxorubicin, and improved efficacy through corrected pharmacokinetic exposure for vincristine, cytarabine, irinotecan, and CPX-351 [15,18,3,8,25,10]. These outcomes rest on a formulation-stability foundation — control of phospholipid hydrolysis and oxidative degradation during manufacture and storage — that is rarely visible in the clinical literature but is a precondition for every trial result summarized here [11,13,34]. For drugs whose principal clinical limitation is chemical or pharmacokinetic instability in free solution, the accumulated evidence supports liposomal encapsulation as a validated, clinically consequential correction rather than a formulation curiosity, while underscoring that its benefit is drug-specific and depends on manufacturing stability control that must be demonstrated for each new candidate rather than assumed from the platform's earlier successes [35,36].

6. References

1. Barenholz, Y. (2012) Doxil - the first FDA-approved nano-drug: lessons learned. *Journal of Controlled Release*, 160(2): 117-134.
2. Gabizon, A., Shmeeda, H., Barenholz, Y. (2003) Pharmacokinetics of pegylated liposomal doxorubicin: review of animal and human studies. *Clinical Pharmacokinetics*, 42(5): 419-436.
3. Silverman, J.A., Deitcher, S.R. (2013) Marqibo (vincristine sulfate liposome injection) improves the pharmacokinetics and pharmacodynamics of vincristine. *Cancer Chemotherapy and Pharmacology*, 71(3): 555-564.
4. Gabizon, A., Papahadjopoulos, D. (1988) Liposome formulations with prolonged circulation time in blood and enhanced uptake by tumors. *Proceedings of the National Academy of Sciences USA*, 85(18): 6949-6953.
5. Allen, T.M., Cullis, P.R. (2013) Liposomal drug delivery systems: from concept to clinical applications. *Advanced Drug Delivery Reviews*, 65(1): 36-48.
6. Adler-Moore, J., Proffitt, R.T. (2002) AmBisome: liposomal formulation, structure, mechanism of action and pre-clinical experience. *Journal of Antimicrobial Chemotherapy*, 49(Suppl 1): 21-30.
7. Barenholz, Y. (2001) Liposome application: problems and prospects. *Current Opinion in Colloid and Interface Science*, 6(1): 66-77.
8. Glantz, M.J., LaFollette, S., Jaeckle, K.A., Shapiro, W., Swinnen, L., Rozental, J.R., et al. (1999) Randomized trial of a slow-release version of cytarabine for the intrathecal treatment of lymphomatous meningitis. *Journal of Clinical Oncology*, 17(10): 3110-3116.
9. Kim, E.S. (2015) Liposomal irinotecan: first global approval. *Drugs*, 75(14): 1663-1669.
10. Lancet, J.E., Uy, G.L., Cortes, J.E., Newell, L.F., Lin, T.L., Ritchie, E.K., et al. (2018) CPX-351 (cytarabine and daunorubicin) liposome for injection versus conventional cytarabine plus daunorubicin in older patients with newly diagnosed secondary acute myeloid leukemia. *Journal of Clinical Oncology*, 36(26): 2684-2692.
11. Grit, M., Crommelin, D.J.A. (1993) Chemical stability of liposomes: implications for their physical stability. *Chemistry and Physics of Lipids*, 64(1-3): 3-18.
12. Crommelin, D.J.A. (1984) Influence of lipid composition and ionic strength on the physical stability of liposomes. *Journal of Pharmaceutical Sciences*, 73(11): 1559-1563.
13. Zuidam, N.J., Crommelin, D.J.A. (1995) Chemical hydrolysis of phospholipids. *Journal of Pharmaceutical Sciences*, 84(9): 1113-1119.
14. Grit, M., de Smidt, J.H., Struijke, A., Crommelin, D.J.A. (1989)

- Hydrolysis of phosphatidylcholine in aqueous liposome dispersions. *International Journal of Pharmaceutics*, 50(1): 1-6.
15. Walsh, T.J., Finberg, R.W., Arndt, C., Hiemenz, J., Schwartz, C., Bodensteiner, D., et al. (1999) Liposomal amphotericin B for empirical therapy in patients with persistent fever and neutropenia. *New England Journal of Medicine*, 340(10): 764-771.
 16. Wingard, J.R., White, M.H., Anaissie, E., Raffalli, J., Goodman, J., Arrieta, A. (2000) A randomized, double-blind comparative trial evaluating the safety of liposomal amphotericin B versus amphotericin B lipid complex in the empirical treatment of febrile neutropenia. *Clinical Infectious Diseases*, 31(5): 1155-1163.
 17. Cornely, O.A., Maertens, J., Bresnik, M., Ebrahimi, R., Ullmann, A.J., Bouza, E., et al. (2007) Liposomal amphotericin B as initial therapy for invasive mold infection: a randomized trial comparing a high-loading dose regimen with standard dosing (AmBiLoad trial). *Clinical Infectious Diseases*, 44(10): 1289-1297.
 18. O'Brien, M.E.R., Wigler, N., Inbar, M., Rosso, R., Grischke, E., Santoro, A., et al. (2004) Reduced cardiotoxicity and comparable efficacy in a phase III trial of pegylated liposomal doxorubicin HCl (CAELYX/Doxil) versus conventional doxorubicin for first-line treatment of metastatic breast cancer. *Annals of Oncology*, 15(3): 440-449.
 19. Batist, G., Ramakrishnan, G., Rao, C.S., Chandrasekharan, A., Gutheil, J., Guthrie, T., et al. (2001) Reduced cardiotoxicity and preserved antitumor efficacy of liposome-encapsulated doxorubicin and cyclophosphamide compared with conventional doxorubicin and cyclophosphamide in a randomized, multicenter trial of metastatic breast cancer. *Journal of Clinical Oncology*, 19(5): 1444-1454.
 20. Harris, L., Batist, G., Belt, R., Rovira, D., Navari, R., Azarnia, N., et al. (2002) Liposome-encapsulated doxorubicin compared with conventional doxorubicin in a randomized multicenter trial as first-line therapy of metastatic breast carcinoma. *Cancer*, 94(1): 25-36.
 21. Rivera, E., Valero, V., Arun, B., Royce, M., Adinin, R., Hoelzer, K., et al. (2002) Phase II study of pegylated liposomal doxorubicin in the treatment of advanced breast cancer. *The Oncologist*, 7(4): 358-361.
 22. Krishna, R., Webb, M.S., St Onge, G., Mayer, L.D. (2001) Liposomal and nonliposomal drug pharmacokinetics after intravenous administration and pharmacodynamics of liposomal doxorubicin in tumor-bearing mice. *Journal of Pharmacology and Experimental Therapeutics*, 298(3): 1206-1212.
 23. O'Brien, S., Schiller, G., Lister, J., Damon, L., Goldberg, S., Aulitzky, W., et al. (2013) High-dose vincristine sulfate liposome injection for advanced, relapsed, and refractory adult Philadelphia chromosome-negative acute lymphoblastic leukemia. *Journal of Clinical Oncology*, 31(6): 676-683.
 24. Bomgaars, L., Geyer, J.R., Franklin, J., Dahl, G., Park, J., Winick, N.J., et al. (2004) Phase I trial of intrathecal liposomal cytarabine in children with neoplastic meningitis. *Journal of Clinical Oncology*, 22(19): 3916-3921.
 25. Wang-Gillam, A., Li, C.P., Bodoky, G., Dean, A., Shan, Y.S., Jameson, G., et al. (2016) Nanoliposomal irinotecan with fluorouracil and folinic acid in metastatic pancreatic cancer after previous gemcitabine-based therapy (NAPOLI-1): a global, randomised, open-label, phase 3 trial. *Lancet*, 387(10018): 545-557.
 26. Mayer, L.D., Tai, L.C., Bally, M.B., Mitlenes, G.N., Ginsberg, R.S., Cullis, P.R. (1989) Characterization of liposomal systems containing doxorubicin entrapped in response to pH gradients. *Biochimica et Biophysica Acta*, 978(2): 246-256.
 27. Amselem, S., Gabizon, A., Barenholz, Y. (1990) A large-scale method for the preparation of sterile and non-pyrogenic

- liposomal formulations of defined size distributions. Journal of Pharmaceutical Sciences, 79(11): 1045-1052.
28. Working, P.K., Dayan, A.D. (1996) Pharmacological-toxicological expert report: CAELYX (Stealth liposomal doxorubicin HCl). Human and Experimental Toxicology, 15(9): 751-785.
29. Sercombe, L., Veerati, T., Moheimani, F., Wu, S.Y., Sood, A.K., Hua, S. (2015) Advances and challenges of liposome assisted drug delivery. Frontiers in Pharmacology, 6: 286.
30. Immordino, M.L., Dosio, F., Cattel, L. (2006) Stealth liposomes: review of the basic science, rationale, and clinical applications, existing and potential. International Journal of Nanomedicine, 1(3): 297-315.
31. Bulbake, U., Doppalapudi, S., Kommineni, N., Khan, W. (2017) Liposomal formulations in clinical use: an updated review. Pharmaceutics, 9(2): 12.
32. Chang, H.I., Yeh, M.K. (2012) Clinical development of liposome-based drugs: formulation, characterization, and therapeutic efficacy. International Journal of Nanomedicine, 7: 49-60.
33. Yingchoncharoen, P., Kalinowski, D.S., Richardson, D.R. (2016) Lipid-based drug delivery systems in cancer therapy: what is available and what is yet to come. Pharmacological Reviews, 68(3): 701-787.
34. Crommelin, D.J.A., van Hoogevest, P., Storm, G. (2020) The role of liposomes in clinical nanomedicine development. What now? Now what? Journal of Controlled Release, 318: 256-263.
35. Cattel, L., Ceruti, M., Dosio, F. (2003) From conventional to stealth liposomes: a new frontier in cancer chemotherapy. Tumori, 89(3): 237-249.
36. Drummond, D.C., Meyer, O., Hong, K., Kirpotin, D.B., Papahadjopoulos, D. (1999) Optimizing liposomes for delivery of chemotherapeutic agents to solid tumors. Pharmacological Reviews, 51(4): 691-743.