

Current Research in Surgery

Review Article

Emerging Role of postoperative complication tracking in the Diagnosis of acute appendicitis

Nolan S^{1*}, Murphy S² and Ryan O³

¹Division of General Surgery, Don Center for Surgical Research, Wales

²Department of Clinical General Surgery, Tajo Institute of Surgical Research, Wales

³Department of General Surgery, Neva Research Center for General Surgery, Scotland

*Address for Correspondence: Siobhan Nolan, Division of General Surgery, Don Center for Surgical Research, Wales; E-mail: siobhan.nolan@don.edu

Received: 28 July 2025; Accepted: 28 October 2025; Published: 2 February 2026.

Citation of this article: Nolan S, Murphy S, Ryan O (2026) Emerging Role of postoperative complication tracking in the Diagnosis of acute appendicitis. Curr Res Surg, 9(1): 01-06. DOI: [10.5281/zenodo.22813470](https://doi.org/10.5281/zenodo.22813470)

Copyright: © 2026 Siobhan Nolan, et al. This is an open access article distributed under the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Diagnostic pathways for acute appendicitis, built on clinical scoring, imaging, and inflammatory biomarkers, remain imperfect, producing both missed and negative diagnoses despite decades of refinement [1,10].

Objective: This review synthesizes evidence on how systematic postoperative complication tracking, developed originally for surgical quality audit, feeds back into and refines the diagnostic process for acute appendicitis.

Methods: We synthesized clinical scoring validation studies, imaging and biomarker accuracy reviews, randomized trials of non-operative management, and national audit and registry reports using structured complication-tracking frameworks such as NSQIP and the Clavien-Dindo classification [12,11].

Findings: Complication data linked to diagnostic pathway intervals show that delayed or equivocal diagnosis is associated with higher rates of surgical site infection, abscess, and reoperation [19,17], negative appendectomy rates have fallen substantially with routine imaging adoption [21], and treatment-failure tracking in antibiotic-first trials reveals imaging-derived predictors of subsequent complication [23,26].

Conclusions: Postoperative complication tracking is emerging as a diagnostic feedback mechanism, not merely a retrospective audit exercise, and closing the loop between pre-operative diagnosis and tracked outcomes should improve appendicitis diagnostic pathways going forward [20,24].

Keywords: acute appendicitis; postoperative complications; diagnostic accuracy; surgical audit; negative appendectomy

1. Introduction

Acute appendicitis remains one of the most common surgical emergencies worldwide, and despite decades of refinement in clinical scoring, imaging, and laboratory testing, the diagnostic pathway is still imperfect at both ends: patients with true appendicitis are still occasionally sent home, and patients without it are still occasionally operated on [1,2]. The Alvarado score, published in 1986, remains the most widely taught bedside instrument for early diagnosis, combining symptoms, signs, and a leukocyte count into a single ordinal scale [3], and later systematic reviews confirmed that it performs reasonably well as a rule-out tool but is less reliable as a stand-alone rule-in instrument, particularly in women and children [4]. The Appendicitis Inflammatory Response score and the Pediatric Appendicitis Score were both developed to address specific shortcomings of the Alvarado instrument, the former by incorporating C-reactive protein and a more graded assessment of peritoneal signs [5], the latter by adapting the approach to a pediatric population with different baseline inflammatory markers and pain descriptions [6].

Imaging has progressively supplemented, and in many centers effectively replaced, purely clinical diagnosis. Meta-analyses of ultrasonography and computed tomography have consistently shown that CT achieves superior sensitivity and specificity to ultrasound for adult appendicitis, at the cost of radiation exposure that has driven interest in low-dose and MRI-based protocols, particularly in children and young adults [7,8]. A Cochrane review of CT for suspected appendicitis in adults confirmed high diagnostic accuracy across pooled cohorts while flagging substantial heterogeneity in scanning protocols and reference standards between studies [9]. Andersson's meta-analysis of clinical and laboratory diagnosis likewise found that no single test or score reliably eliminates diagnostic uncertainty, reinforcing that most modern pathways rely on a combination of scoring, imaging, and observation [10].

What has received comparatively less attention is the diagnostic information embedded in what happens after the operation. Every appendectomy generates a postoperative course, a pathology report, and, increasingly, a structured complication record, whether through national audit programs, institutional registries, or classification systems such as the Clavien-Dindo grading scale [11]. This review examines the evidence that systematic tracking of postoperative complications, rather than functioning only as a quality metric, feeds back into and refines the diagnostic process for acute appendicitis, by exposing missed or delayed diagnoses, quantifying the cost of diagnostic uncertainty, and identifying subgroups in whom the pre-operative diagnostic pathway underperforms.

2. Methods

We synthesized the peer-reviewed literature on the diagnosis and postoperative management of acute appendicitis, drawing on clinical scoring system validation studies, imaging accuracy meta-analyses, large administrative and audit-based outcome cohorts, randomized trials of non-operative management, and reports describing structured surgical complication tracking systems such as the National Surgical Quality Improvement Program (NSQIP) and the Clavien-Dindo classification [12,11]. Sources were identified from major general surgery, emergency medicine, and gastroenterology journals and from established international guideline documents, including the World Society of Emergency Surgery Jerusalem guidelines and the European Association for Endoscopic Surgery consensus statement [13,14]. Studies were included where they reported diagnostic performance of a clinical, laboratory, or imaging test for appendicitis; described postoperative complication rates linked to diagnostic timing, negative appendectomy, or perforation status; or described the structure and function of a complication-tracking system applicable to appendectomy outcomes. Findings were organized

thematically around three questions: how diagnostic accuracy is currently measured, how postoperative complication data are captured, and how the two domains inform one another in practice.

3. Results

Diagnostic performance of clinical scoring systems

Across validation cohorts, the Alvarado score at the conventional cut-off performs with moderate sensitivity and specificity, prompting numerous efforts at modification [3,4]. The Appendicitis Inflammatory Response score, validated head-to-head against Alvarado, achieved a lower rate of indeterminate classifications and correspondingly reduced the proportion of patients requiring further imaging before a management decision could be made [5]. Population-based incidence data show that the overall rate of appendicitis has been relatively stable over recent decades even as the mix of imaging-confirmed versus purely clinical diagnoses has shifted substantially, a trend attributable to the wider availability of cross-sectional imaging rather than to any true change in disease incidence [15,2].

Imaging, biomarkers, and diagnostic delay

Diagnostic biomarkers, principally C-reactive protein, white blood cell count, and more recently procalcitonin, have been evaluated extensively as adjuncts to clinical scoring. A meta-analysis of these three markers found that no single biomarker reliably distinguished complicated from uncomplicated appendicitis, though combinations modestly improved discrimination over any marker alone [16]. Delay from symptom onset to operative intervention correlates directly with perforation risk: cohort data quantifying "the risk of rupture" as a function of elapsed time found a measurable and continuous increase in perforation probability with each additional hour of untreated appendicitis beyond the first day of symptoms [17], a relationship consistent with the broader observation that non-perforated and perforated appendicitis behave almost as epidemiologically distinct

entities rather than points on a single severity continuum [18].

Table 1: Reported diagnostic performance characteristics of principal clinical scoring systems for acute appendicitis [3,5,6,4].

Scoring system	Population	Key components	Reported performance
Alvarado score	Adults	Symptoms, signs, leukocytosis	Moderate sensitivity; limited specificity at rule-in threshold
Appendicitis Inflammatory Response score	Adults	Peritoneal signs, CRP, leukocytosis	Fewer indeterminate classifications than Alvarado
Pediatric Appendicitis Score	Children	Symptoms, signs, leukocytosis	Comparable accuracy adapted to pediatric presentation

Postoperative complication tracking as a diagnostic signal

Structured complication tracking, whether through NSQIP-style risk-adjusted audit or through simpler grading systems such as Clavien-Dindo, converts an individual patient's postoperative course into a comparable, quantifiable outcome [12,11]. When these outcomes are linked back to pre-operative diagnostic pathway data, several patterns emerge consistently across national audits: patients who underwent surgery after a longer diagnostic interval, or after an equivocal imaging study, show higher rates of surgical site infection, intra-abdominal abscess, and reoperation than patients diagnosed promptly [19,20]. Negative appendectomy, once accepted as an unavoidable trade-off for avoiding perforation, is itself captured by complication-tracking frameworks as an adverse outcome, and its documented rate has fallen substantially in centers that adopted routine pre-operative imaging, a decline directly attributable to the diagnostic pathway rather than to any change in surgical technique [21,22].

Table 1 summarizes reported diagnostic performance characteristics of the principal clinical scoring systems discussed above, and Figure 1 illustrates the pattern of

falling negative appendectomy rates alongside increasing routine imaging use reported across successive audit periods.

Table 2: Thirty-day complication rates stratified by perforation status, drawn from national audit and registry cohorts [19,20,29].			
Outcome	Non-perforated	Perforated	Tracking source
Surgical site infection	Low	Substantially higher	National audit / NSQIP-style registry
Intra-abdominal abscess	Rare	Meaningfully elevated	National audit / registry
Reoperation	Rare	Elevated	National audit / registry
Length of stay	Short	Markedly prolonged	Administrative database




Figure 1: Negative appendectomy rate by audit period, falling from approximately 18% before routine pre-operative imaging to approximately 5% after routine CT/ultrasound adoption, a decline attributed to imaging-informed diagnostic pathways rather than to changes in operative technique [21,22].




Figure 2: Thirty-day postoperative complication incidence by diagnostic-to-operative interval, rising from approximately 4% when surgery followed diagnosis within 12 hours to approximately 22% beyond 48 hours, consistent with time-dependent perforation risk [17,19].

Randomized trials of non-operative, antibiotic-first management add a further dimension to complication

tracking, since the complication profile of the antibiotic arm, including treatment failure requiring delayed appendectomy, becomes itself a diagnostic and prognostic signal about which patients were correctly identified as having genuinely uncomplicated disease [23,24,25]. Meta-analyses pooling these trials found that a meaningful minority of patients randomized to antibiotics ultimately required appendectomy within the following year, and treatment failure was more common in patients whose index imaging had shown a fecalith, suggesting that imaging-derived features captured at diagnosis predict a complication that only becomes apparent during follow-up [26,27]. Table 2 sets out complication rates stratified by perforation status drawn from these tracking-based cohorts, and Figure 2 shows thirty-day complication incidence stratified by diagnostic pathway interval.

4. Discussion

Complication registries and quality improvement

The Department of Veterans Affairs' NSQIP program was explicitly designed to make postoperative outcomes risk-adjusted and comparable across institutions, and its methodology has since been adapted broadly across general surgery, including appendectomy-specific outcome reporting [12]. Hospital-level quality metrics derived from such registries correlate with resource use and, in some analyses, with overall cost of inpatient surgical care, suggesting that complication tracking captures information about diagnostic and perioperative process quality that is not visible from operative volume or case mix alone [28]. The laparoscopic-versus-open comparison literature illustrates the same principle in miniature: large administrative-database analyses found that laparoscopic appendectomy was associated with lower rates of wound infection but a higher rate of intra-abdominal abscess than open appendectomy, a pattern only detectable because outcomes were tracked systematically across a sufficiently large cohort, and later

Cochrane synthesis confirmed the finding while noting persistent heterogeneity in how "complication" was defined across included trials [29,30].

Limits of complication tracking as a diagnostic tool

Complication tracking is not without important limitations when used as a diagnostic feedback mechanism. Clavien-Dindo grading, while reproducible, was designed as a general surgical severity taxonomy rather than one specific to appendicitis, and it does not by itself distinguish a complication attributable to diagnostic delay from one attributable to unrelated patient comorbidity [11]. Registries such as the Prospective Observational Study on Acute Appendicitis Worldwide highlight substantial international variation in both diagnostic pathway and complication definitions, which complicates direct comparison of tracked outcomes across health systems [20]. The management of the periappendicular abscess and interval appendectomy question illustrates this well: randomized comparison of interval appendectomy against imaging follow-up found a measurable, if modest, rate of missed appendiceal neoplasm in the follow-up arm, a diagnostic miss that only complication and pathology tracking over the following year was able to surface [31]. Guideline bodies have increasingly recommended standardized reporting of exactly this kind of delayed diagnostic information as part of routine appendicitis audit [13,14,32].

Diagnostic accuracy for suspected acute appendicitis in the emergency department setting remains an active area of evidence synthesis, and contemporary evidence-based reviews increasingly frame complication and reoperation rates, rather than sensitivity and specificity alone, as outcome measures that should inform which diagnostic pathway a given institution adopts [33,34]. This reframing treats the diagnostic pathway and the postoperative complication record as a single continuous audit loop rather than as two separate stages of care, an approach consistent with the broader shift toward outcome-based,

risk-adjusted surgical quality measurement pioneered by programs such as NSQIP [12,28].

5. Conclusion

Postoperative complication tracking, developed originally as a quality-improvement and audit tool, has an increasingly well-documented secondary role as a diagnostic feedback mechanism for acute appendicitis. By linking structured outcome data, including surgical site infection, abscess formation, reoperation, and missed pathology, back to the pre-operative diagnostic pathway, institutions and registries have been able to identify where clinical scoring, imaging, and biomarker-based diagnosis underperform, and to quantify the real clinical cost of both diagnostic delay and negative appendectomy [11,19,21]. As non-operative management strategies expand the range of diagnostic-therapeutic decisions clinicians must make in real time, complication tracking is positioned to become not merely a retrospective audit exercise but an active, closed-loop component of the diagnostic process itself [23,24,20].

6. References

1. Bhangu, A., Soreide, K., Di Saverio, S., Assarsson, J.H., Drake, F.T. (2015) Acute appendicitis: modern understanding of pathogenesis, diagnosis, and management. *Lancet*, 386(10000): 1278-1287. DOI: [10.1016/S0140-6736\(15\)00275-5](https://doi.org/10.1016/S0140-6736(15)00275-5)
2. Ferris, M., Quan, S., Kaplan, B.S., Molodecky, N., Ball, C.G., Chernoff, G.W., et al. (2017) The global incidence of appendicitis: a systematic review of population-based studies. *Annals of Surgery*, 266(2): 237-241. DOI: [10.1097/SLA.0000000000002188](https://doi.org/10.1097/SLA.0000000000002188)
3. Alvarado, A. (1986) A practical score for the early diagnosis of acute appendicitis. *Annals of Emergency Medicine*, 15(5): 557-564. DOI: [10.1016/S0196-0644\(86\)80993-3](https://doi.org/10.1016/S0196-0644(86)80993-3)
4. Ohle, R., O'Reilly, F., O'Brien, K.K., Fahey, T., Dimitrov, B.D. (2011) The Alvarado score for predicting acute appendicitis: a systematic review. *BMC Medicine*, 9: 139. DOI: [10.1186/1745-6215-9-139](https://doi.org/10.1186/1745-6215-9-139)

[10.1186/1741-7015-9-139](https://doi.org/10.1186/1741-7015-9-139)

5. Andersson, M., Andersson, R.E. (2008) The Appendicitis Inflammatory Response score: a tool for the diagnosis of acute appendicitis that outperforms the Alvarado score. *World Journal of Surgery*, 32(8): 1843-1849. DOI: [10.1007/s00268-008-9649-y](https://doi.org/10.1007/s00268-008-9649-y)
6. Samuel, M. (2002) Pediatric appendicitis score. *Journal of Pediatric Surgery*, 37(6): 877-881. DOI: [10.1053/jpsu.2002.32893](https://doi.org/10.1053/jpsu.2002.32893)
7. Terasawa, T., Blackmore, C.C., Bent, S., Kohlwes, R.J. (2004) Systematic review: computed tomography and ultrasonography to detect acute appendicitis in adults and adolescents. *Annals of Internal Medicine*, 141(7): 537-546. DOI: [10.7326/0003-4819-141-7-200410050-00011](https://doi.org/10.7326/0003-4819-141-7-200410050-00011)
8. Kim, K., Kim, Y.H., Kim, S.Y., Kim, S., Lee, Y.J., Kim, K.P., et al. (2012) Low-dose radiation CT for the diagnosis of appendicitis in adolescents and young adults. *New England Journal of Medicine*, 366(17): 1596-1605. DOI: [10.1056/NEJMoa1110734](https://doi.org/10.1056/NEJMoa1110734)
9. Rud, B., Vejborg, T.S., Rapoport, E.D., Reitsma, J.B., Wille-Jorgensen, P. (2019) Computed tomography for diagnosis of acute appendicitis in adults. *Cochrane Database of Systematic Reviews*, 2019(11): CD009977. DOI: [10.1002/14651858.CD009977.pub2](https://doi.org/10.1002/14651858.CD009977.pub2)
10. Andersson, R.E. (2004) Meta-analysis of the clinical and laboratory diagnosis of appendicitis. *British Journal of Surgery*, 91(1): 28-37. DOI: [10.1002/bjs.4464](https://doi.org/10.1002/bjs.4464)
11. Dindo, D., Demartines, N., Clavien, P.A. (2004) Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Annals of Surgery*, 240(2): 205-213. DOI: [10.1097/01.sla.0000133083.54934.ae](https://doi.org/10.1097/01.sla.0000133083.54934.ae)
12. Khuri, S.F., Daley, J., Henderson, W., Hur, K., Demakis, J., Aust, J.B., et al. (1998) The Department of Veterans Affairs' NSQIP: the first national, validated, outcome-based, risk-adjusted, and peer-controlled program for the measurement and enhancement of the quality of surgical care. *Annals of Surgery*, 228(4): 491-507. DOI: [10.1097/0000658-199810000-00006](https://doi.org/10.1097/0000658-199810000-00006)
13. Di Saverio, S., Podda, M., De Simone, B., Ceresoli, M., Augustin, G., Gori, A., et al. (2020) Diagnosis and treatment of acute appendicitis: 2020 update of the WSES Jerusalem guidelines. *World Journal of Emergency Surgery*, 15: 27. DOI: [10.1186/s13017-020-00306-3](https://doi.org/10.1186/s13017-020-00306-3)
14. Gorter, R.R., Eker, H.H., Gorter-Stam, M.A., Abis, G.S., Acharya, A., Ankersmit, M., et al. (2016) Diagnosis and management of acute appendicitis. EAES consensus development conference 2015. *Surgical Endoscopy*, 30(11): 4668-4690. DOI: [10.1007/s00464-016-5245-7](https://doi.org/10.1007/s00464-016-5245-7)
15. Coward, S., Kareemi, H., Clement, F., Zimmer, S., Dixon, E., Ball, C.G., et al. (2016) Incidence of appendicitis over time: a comparative analysis of an administrative healthcare database and a pathology-proven appendicitis registry. *PLoS ONE*, 11(11): e0165161. DOI: [10.1371/journal.pone.0165161](https://doi.org/10.1371/journal.pone.0165161)
16. Yu, C.W., Juan, L.I., Wu, M.H., Shen, C.J., Wu, J.Y., Lee, C.C. (2013) Systematic review and meta-analysis of the diagnostic accuracy of procalcitonin, C-reactive protein and white blood cell count for suspected acute appendicitis. *British Journal of Surgery*, 100(3): 322-329. DOI: [10.1002/bjs.9008](https://doi.org/10.1002/bjs.9008)
17. Bickell, N.A., Aufses, A.H., Rojas, M., Bodian, C. (2006) How time affects the risk of rupture in appendicitis. *Journal of the American College of Surgeons*, 202(3): 401-406. DOI: [10.1016/j.jamcollsurg.2005.11.016](https://doi.org/10.1016/j.jamcollsurg.2005.11.016)
18. Livingston, E.H., Woodward, W.A., Sarosi, G.A., Haley, R.W. (2007) Disconnect between incidence of nonperforated and perforated appendicitis: implications for pathophysiology and management. *Annals of Surgery*, 245(6): 886-892. DOI: [10.1097/01.sla.0000256391.05233.aa](https://doi.org/10.1097/01.sla.0000256391.05233.aa)
19. van Rossem, C.C., Bolmers, M.D., Schreinemacher, M.H., van Geloven, A.A., Bemelman, W.A. (2016) Prospective

- nationwide outcome audit of surgery for suspected acute appendicitis. *British Journal of Surgery*, 103(1): 144-151. DOI: [10.1002/bjs.9964](https://doi.org/10.1002/bjs.9964)
20. Sartelli, M., Baiocchi, G.L., Di Saverio, S., Ferrara, F., Labricciosa, F.M., Ansaloni, L., et al. (2018) Prospective Observational Study on acute Appendicitis Worldwide (POSAW). *World Journal of Emergency Surgery*, 13: 19. DOI: [10.1186/s13017-018-0179-0](https://doi.org/10.1186/s13017-018-0179-0)
 21. Drake, F.T., Flum, D.R. (2013) Improvement in the diagnosis of appendicitis. *Advances in Surgery*, 47: 299-328. DOI: [10.1016/j.yasu.2013.03.001](https://doi.org/10.1016/j.yasu.2013.03.001)
 22. Moris, D., Paulson, E.K., Pappas, T.N. (2021) Diagnosis and management of acute appendicitis in adults: a review. *JAMA*, 326(22): 2299-2311. DOI: [10.1001/jama.2021.20502](https://doi.org/10.1001/jama.2021.20502)
 23. Salminen, P., Paajanen, H., Rautio, T., Nordstrom, P., Aarnio, M., Rantanen, T., et al. (2015) Antibiotic therapy vs appendectomy for treatment of uncomplicated acute appendicitis: the APPAC randomized clinical trial. *JAMA*, 313(23): 2340-2348. DOI: [10.1001/jama.2015.6154](https://doi.org/10.1001/jama.2015.6154)
 24. CODA Collaborative (2020) A randomized trial comparing antibiotics with appendectomy for appendicitis. *New England Journal of Medicine*, 383(20): 1907-1919. DOI: [10.1056/NEJMoa2014320](https://doi.org/10.1056/NEJMoa2014320)
 25. Sippola, S., Haijanen, J., Gronroos, J., Rautio, T., Nordstrom, P., Rantanen, T., et al. (2021) Effect of oral moxifloxacin vs intravenous ertapenem plus oral levofloxacin for treatment of uncomplicated acute appendicitis: the APPAC II randomized clinical trial. *JAMA*, 325(4): 353-362. DOI: [10.1001/jama.2020.23525](https://doi.org/10.1001/jama.2020.23525)
 26. Sallinen, V., Akl, E.A., You, J.J., Agarwal, A., Shoucair, S., Vandvik, P.O., et al. (2016) Meta-analysis of antibiotics versus appendicectomy for non-perforated acute appendicitis. *British Journal of Surgery*, 103(6): 656-667. DOI: [10.1002/bjs.10147](https://doi.org/10.1002/bjs.10147)
 27. Poprom, N., Wilasrusmee, C., Attia, J., McEvoy, M., Thakkestian, A., Rattanasiri, S. (2019) The efficacy of antibiotic treatment versus surgical treatment of uncomplicated acute appendicitis: systematic review and meta-analysis. *American Journal of Surgery*, 218(1): 192-200. DOI: [10.1016/j.amjsurg.2018.10.009](https://doi.org/10.1016/j.amjsurg.2018.10.009)
 28. Birkmeyer, J.D., Gust, C., Dimick, J.B., Birkmeyer, N.J., Skinner, J.S. (2012) Hospital quality and the cost of inpatient surgery in the United States. *Annals of Surgery*, 255(1): 1-5. DOI: [10.1097/SLA.0b013e3182402c17](https://doi.org/10.1097/SLA.0b013e3182402c17)
 29. Guller, U., Hervey, S., Purves, H., Muhlbaier, L.H., Peterson, E.D., Eubanks, S., et al. (2004) Laparoscopic versus open appendectomy: outcomes comparison based on a large administrative database. *Annals of Surgery*, 239(1): 43-52. DOI: [10.1097/01.sla.0000103071.35986.c1](https://doi.org/10.1097/01.sla.0000103071.35986.c1)
 30. Sauerland, S., Jaschinski, T., Neugebauer, E.A. (2010) Laparoscopic versus open surgery for suspected appendicitis. *Cochrane Database of Systematic Reviews*, 2010(10): CD001546. DOI: [10.1002/14651858.CD001546.pub3](https://doi.org/10.1002/14651858.CD001546.pub3)
 31. Mallinen, J., Rautio, T., Gronroos, J., Rantanen, T., Nordstrom, P., Savolainen, H., et al. (2019) Risk of appendiceal neoplasm in periappendicular abscess in patients treated with interval appendectomy vs follow-up with CT or MRI: 1-year outcomes of the multicenter, randomized APPAC III trial. *JAMA Surgery*, 154(3): 200-207. DOI: [10.1001/jamasurg.2018.4373](https://doi.org/10.1001/jamasurg.2018.4373)
 32. Wagner, M., Tubre, D.J., Asensio, J.A. (2018) Evolution and current trends in the management of acute appendicitis. *Surgical Clinics of North America*, 98(5): 1005-1023. DOI: [10.1016/j.suc.2018.05.006](https://doi.org/10.1016/j.suc.2018.05.006)
 33. Shogilev, D.J., Duus, N., Odom, S.R., Shapiro, N.I. (2014) Diagnosing appendicitis: evidence-based review of the diagnostic approach in 2014. *Western Journal of Emergency Medicine*, 15(7): 859-871. DOI: [10.5811/westjem.2014.9.21568](https://doi.org/10.5811/westjem.2014.9.21568)
 34. Flum, D.R. (2015) Acute appendicitis - appendectomy or the "antibiotics first" strategy. *New England Journal of Medicine*, 372(20): 1937-1943. DOI: [10.1056/NEJMcp1215006](https://doi.org/10.1056/NEJMcp1215006)