Somato Publications

ISSN: XXXX-XXXX

International Journal of Arthritis

Review Article

Clinical Outcomes of biologic disease-modifying therapy in Patients with rheumatoid arthritis

Diallo C[1*], Mensah C[2] and Banda K[3]

[1]Department of Arthritis Research, Murray Institute of Rheumatology, Zimbabwe

[2]Department of Rheumatology, Guadalquivir Research Center for Rheumatology, South Africa

[3]Division of Arthritis Research, Snake Academy of Arthritis Research, South Africa

*Address for Correspondence: Chiamaka Diallo, Department of Arthritis Research, Murray Institute of Rheumatology, Zimbabwe; E-mail: chiamaka.diallo@murray.edu

Received: 19 May 2022; Accepted: 25 August 2022; Published: 15 November 2022.

Citation of this article: Diallo C, Mensah C, Banda K (2022) Clinical Outcomes of biologic disease-modifying therapy in Patients with rheumatoid arthritis. Int J Arthritis, 4(1): 01-06. DOI: 10.5281/zenodo.22813981

Copyright: © 2022 Chiamaka Diallo, et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Biologic disease-modifying antirheumatic drugs (bDMARDs) transformed the management of rheumatoid arthritis (RA) by delivering sustained clinical response and structural protection unattainable with conventional synthetic agents alone.

Objective: This review synthesizes the clinical trial and registry evidence on outcomes with biologic DMARD therapy in RA, covering efficacy across drug classes, predictors of response, comparative and combination strategies, and infection risk.

Methods: We synthesized pivotal randomized controlled trials, long-term extension studies, and large national biologics registries addressing TNF-inhibitor and non-TNF biologic efficacy, treatment sequencing, and safety outcomes in RA.

Findings: TNF inhibitors combined with methotrexate produce markedly higher ACR response rates and near-complete arrest of radiographic progression compared with methotrexate alone, non-TNF mechanisms targeting B cells, T-cell costimulation, and interleukin-6 retain efficacy after anti-TNF failure, and serious infection risk, while real and concentrated in the first six months of treatment, is broadly comparable across biologic classes.

Conclusions: Biologic DMARD therapy, introduced early and used within a treat-to-target strategy aimed at sustained remission, delivers substantial and durable gains in disease control and function, with individualized agent selection and sequencing supported by identifiable predictors of response and by evidence favoring mechanism-switching after anti-TNF failure.

Keywords: rheumatoid arthritis; biologic DMARD; TNF inhibitor; clinical outcomes; serious infection risk

1. Introduction

Rheumatoid arthritis (RA) is a chronic, symmetric, erosive polyarthritis that, left inadequately controlled, produces progressive joint destruction, functional disability, and excess cardiovascular and all-cause mortality. The classification framework jointly issued by the American College of Rheumatology and the European League Against Rheumatism formalized how patients are identified for treatment before irreversible damage accumulates [1], and disease activity is now tracked with composite indices such as the 28-joint Disease Activity Score, which combines joint counts, an acute-phase reactant, and a patient global assessment into a single number that drives treat-to-target decisions in trials and practice [2]. Before biologic disease-modifying antirheumatic drugs (bDMARDs), conventional synthetic agents, principally methotrexate, controlled symptoms in a minority of patients but left many with ongoing synovitis and accumulating disability, as documented in longitudinal cohorts linking joint damage directly to loss of function [3].

The arrival of tumor necrosis factor (TNF) inhibitors in the late 1990s reset expectations for pharmacologic control of RA. Infliximab combined with methotrexate produced sustained clinical and radiographic benefit unattainable with methotrexate alone in the pivotal ATTRACT trial [4], building on an earlier open-label study by Maini and colleagues showing that repeated infliximab infusions produced measurable clinical improvement over background methotrexate [5]. Etanercept demonstrated a comparable effect in the trial reported by Weinblatt and colleagues [6], and adalimumab extended the class with a fully human monoclonal antibody producing significant radiographic, clinical, and functional benefit over methotrexate alone [7]. These agents, followed within a decade by non-TNF biologics targeting B cells, T-cell costimulation, and interleukin-6, transformed RA from a disease managed toward symptomatic tolerability into one managed toward remission, a shift codified in successive EULAR and ACR recommendations [8,9,10,11]. This review synthesizes trial and registry evidence on outcomes with biologic DMARD therapy in RA, covering efficacy across drug classes, predictors of response, combination strategy, and the infection-risk trade-off that accompanies sustained biologic exposure.

2. TNF-inhibitor efficacy: the foundational evidence

The TNF-inhibitor class remains the most extensively studied biologic mechanism in RA, and its founding trials established the methodology, ACR response criteria and radiographic scoring, by which every subsequent biologic has since been judged. In the ATTRACT trial, infliximab added to methotrexate produced significantly higher ACR20, ACR50, and ACR70 response rates than methotrexate alone at week 54, and patients receiving infliximab showed essentially no progression in radiographic score, while the methotrexate-alone group continued to accrue joint damage [4]. This dissociation between clinical response and structural protection became a defining feature distinguishing biologic from conventional synthetic therapy.

Etanercept's registration program showed a parallel pattern. Weinblatt and colleagues demonstrated that etanercept added to background methotrexate produced rapid, sustained ACR20/50/70 responses emerging within two weeks and maintained through six months [6], and follow-up by Genovese and colleagues in early RA confirmed durable clinical, functional, and radiographic benefit extending out several years [12]. Klareskog and colleagues showed in the TEMPO trial that etanercept combined with methotrexate outperformed either drug alone on every domain measured, the clearest head-to-head evidence that combination therapy delivers the greatest structural protection [13]. Adalimumab's PREMIER trial extended this combination-superiority finding to early, aggressive RA, with adalimumab plus methotrexate arresting radiographic progression in far

© 2022 Somato Publications. All rights reserved.

more patients than either monotherapy arm over two years [14], while Keystone and colleagues confirmed sustained ACR response alongside minimal radiographic progression through one year in an independent cohort [7].




Figure 1: ACR20, ACR50, and ACR70 response rates at week 54 in the ATTRACT trial, comparing methotrexate alone with infliximab plus methotrexate, illustrating the magnitude of clinical benefit attributable to TNF-inhibitor combination therapy [4].

3. Non-TNF biologic mechanisms

Mechanistic diversification beyond TNF blockade addressed the substantial minority of patients who fail to respond, or lose response over time, to anti-TNF therapy. Rituximab, a chimeric monoclonal antibody depleting CD20-positive B cells, produced significant ACR response in methotrexate-inadequate-responder patients in the dose-ranging trial by Emery and colleagues [15], and Cohen and colleagues demonstrated that rituximab retained substantial efficacy specifically in patients who had already failed anti-TNF agents, establishing B-cell depletion as a viable second-line mechanism [16]. Abatacept, which blocks T-cell costimulation by binding CD80/CD86, showed comparable efficacy in the methotrexate-inadequate-responder population studied by Kremer and colleagues [17], and Genovese and colleagues confirmed abatacept remained effective in patients refractory to TNF-alpha inhibition, showing that a mechanistically distinct biologic can rescue response after anti-TNF failure [18].

Interleukin-6 receptor blockade with tocilizumab added a third non-TNF mechanism with a distinctive profile. In the AMBITION trial, tocilizumab monotherapy outperformed methotrexate monotherapy, notable because most other biologics show their strongest effect in combination with methotrexate [19]. Emery and colleagues extended this evidence to anti-TNF-refractory patients, showing significant improvement even in a population that had already failed biologic therapy targeting a different pathway [20]. Gottenberg and colleagues directly compared switching to a non-TNF biologic against cycling to a second anti-TNF agent after inadequate response to a first TNF inhibitor, and found switching mechanistic class produced a significantly higher proportion of patients achieving low disease activity, evidence that has shaped sequencing decisions [21].

Table 1: Non-TNF-targeted biologic mechanisms in rheumatoid arthritis and their pivotal trial evidence [15,16,17,18,19,20].

Agent	Target	Pivotal population	Key finding
Rituximab	CD20 (B cells)	MTX-inadequate responders / anti-TNF failures	Significant ACR response and efficacy retained after anti-TNF failure [15,16]
Abatacept	CD80/CD86 costimulation	MTX-inadequate responders / anti-TNF failures	Efficacy in MTX failures and in patients refractory to TNF blockade [17,18]
Tocilizumab	IL-6 receptor	MTX-naive and anti-TNF-refractory	Superior to MTX monotherapy; effective after anti-TNF failure [19,20]

4. Later-generation TNF inhibitors and combination strategy

Certolizumab pegol, a PEGylated Fc-free anti-TNF fragment introduced after the first wave of TNF inhibitors, produced significantly greater clinical response than placebo plus methotrexate in the fifty-two-week RAPID 1 trial reported by Keystone and colleagues, reproducing the core class efficacy signal with a differing dosing

© 2022 Somato Publications. All rights reserved.

schedule [22]. Combination strategy, rather than biologic choice alone, has emerged as a central determinant of long-term outcome. Markusse and colleagues, reporting ten-year outcomes from a tight-control treatment trial in recent-onset RA, found that patients managed with protocolized treat-to-target strategies, several incorporating biologic step-up, achieved sustained low disease activity and preserved function at a decade regardless of which initial strategy arm they were randomized to, underscoring that the target-driven treatment philosophy matters as much as the specific agent [23]. Radner and colleagues quantified the clinical and economic case for this approach, showing that patients achieving formal remission by ACR/EULAR criteria had meaningfully better patient-reported outcomes and lower downstream costs than those achieving only low disease activity [24]. The ACR/EULAR provisional remission definition itself, requiring near-total absence of tender and swollen joints alongside a near-normal acute-phase reactant, was developed specifically to give trials of biologic and combination strategies a stringent, reproducible endpoint [25].

5. Predictors of response and treatment sequencing

Not all patients respond equally to a given biologic, and identifying predictors of response, or of the need to switch mechanism, has been a persistent focus of registry-based research. Finckh and colleagues, analyzing a Swiss biologics registry, found evidence for genuine acquired drug resistance across sequential anti-TNF agents, so that failure of a first anti-TNF drug predicted a measurably lower probability of response to a second agent within the same class [26]. Hyrich and colleagues, using the British Society for Rheumatology Biologics Register, identified baseline predictors of anti-TNF response including lower baseline disability score, male sex, and shorter disease duration, findings that have informed patient selection and expectation-setting [27]. Hetland and colleagues, comparing adalimumab, etanercept, and infliximab head-to-head in a national Danish registry, found broadly similar remission rates and drug adherence across the three agents, with modest differences favoring etanercept and adalimumab over infliximab on retention [28]. Taken together with the sequencing evidence from Finckh, Hyrich, and Gottenberg, these results support a management approach in which biologic choice and sequencing are informed by baseline patient characteristics and early response trajectory rather than applied uniformly [26,27,21].

Table 2: Registry-identified predictors of anti-TNF response and sequencing outcomes in rheumatoid arthritis [26,27,28,21].

Factor / strategy	Source	Association with outcome
Lower baseline disability, male sex, shorter disease duration	British Society for Rheumatology Biologics Register	Predict better anti-TNF response [27]
Failure of first anti-TNF agent	Swiss biologics registry	Predicts reduced response to a second anti-TNF agent [26]
Adalimumab, etanercept, infliximab (head-to-head)	Danish national registry	Broadly similar remission rates; modest retention advantage for etanercept/adalimumab [28]
Switch to non-TNF biologic vs cycle to second anti-TNF	Randomized comparison	Switching mechanism yields higher low-disease-activity rate [21]

6. Infection risk and other safety considerations

The clinical benefit of biologic DMARDs has always been weighed against an increased risk of serious infection, characterized across trial-level meta-analyses and large observational registries. Bongartz and colleagues' meta-analysis of anti-TNF antibody trials found a significantly increased odds of serious infection and a signal for malignancy compared with placebo, a finding that prompted regulatory boxed warnings and closer post-marketing surveillance across the class [29]. Listing and colleagues, using data from the German biologics register, quantified this risk directly in routine practice, finding a significantly elevated rate of serious infection in anti-TNF-treated patients compared with those on conventional synthetic DMARDs, particularly with higher baseline disability and glucocorticoid co-therapy [30]. Galloway and colleagues, analyzing the British registry,

© 2022 Somato Publications. All rights reserved.

refined the temporal pattern of this risk, showing that the excess of serious infection associated with anti-TNF therapy was concentrated in the first six months of treatment, after which the incremental risk relative to conventional therapy narrowed considerably [31].




Figure 2: Serious infection incidence by time since anti-TNF initiation in registry cohorts, showing the excess risk concentrated in the first six months of treatment before narrowing thereafter [31,30].

Singh and colleagues' Cochrane network meta-analysis and overview of biologic adverse effects synthesized this evidence across drug classes, TNF inhibitors, rituximab, abatacept, and tocilizumab alike, and concluded that serious infection risk was elevated to a broadly similar degree across mechanisms, with malignancy and tuberculosis reactivation as additional considerations that vary by agent and baseline screening practice [32]. Current ACR and EULAR guidance incorporates this evidence directly into prescribing recommendations, calling for tuberculosis and hepatitis screening before biologic initiation and individualized risk-benefit discussion in patients with a history of serious infection or malignancy [9,11].

7. Conclusion

Three decades of trial and registry evidence now support a coherent picture of biologic DMARD therapy in rheumatoid arthritis: TNF inhibitors, and the non-TNF mechanisms that followed, reliably produce clinical response and structural protection unattainable with conventional synthetic DMARDs alone, particularly when combined with methotrexate and introduced early before irreversible joint damage accumulates [4,13,14]. Response is not uniform, genuine acquired resistance across sequential agents within a class, and identifiable baseline predictors of response, mean that treatment sequencing benefits from individualization, and switching to a mechanistically distinct biologic after a first anti-TNF failure now has direct comparative evidence behind it [26,27,21]. That benefit carries a real, well-quantified increase in serious infection risk, concentrated in the early months of treatment and broadly comparable across drug classes, that must be weighed against the substantial gains in disease control, function, and quality of life that biologic therapy, used within a treat-to-target strategy aimed at sustained remission, has delivered for a disease that previously left many patients with progressive, disabling joint destruction [29,31,24]. Ongoing registry surveillance, alongside continued refinement of predictors that guide agent selection and sequencing, remains essential to sustaining this balance as the biologic armamentarium continues to expand.

8. References

1. Aletaha, D., Neogi, T., Silman, A.J., Funovits, J., Felson, D.T., Bingham, C.O., et al. (2010) 2010 rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. Arthritis & Rheumatism, 62(9): 2569-2581. DOI: 10.1002/art.27584

2. Prevoo, M.L.L., van 't Hof, M.A., Kuper, H.H., van Leeuwen, M.A., van de Putte, L.B.A., van Riel, P.L.C.M. (1995) Modified disease activity scores that include twenty-eight-joint counts. Arthritis & Rheumatism, 38(1): 44-48. DOI: 10.1002/art.1780380107

3. Scott, D.L., Pugner, K., Kaarela, K., Doyle, D.V., Woolf, A., Holmes, J., et al. (2000) The links between joint damage and disability in rheumatoid arthritis. Rheumatology, 39(2): 122-132. DOI: 10.1093/rheumatology/39.2.122

4. Lipsky, P.E., van der Heijde, D.M., St Clair, E.W., Furst, D.E., Breedveld, F.C., Kalden, J.R., et al. (2000) Infliximab and

© 2022 Somato Publications. All rights reserved.

methotrexate in the treatment of rheumatoid arthritis. New England Journal of Medicine, 343(22): 1594-1602. DOI: 10.1056/NEJM200011303432202

5. Maini, R.N., Breedveld, F.C., Kalden, J.R., Smolen, J.S., Davis, D., Macfarlane, J.D., et al. (1998) Therapeutic efficacy of multiple intravenous infusions of anti-tumor necrosis factor alpha monoclonal antibody combined with low-dose weekly methotrexate in rheumatoid arthritis. Arthritis & Rheumatism, 41(9): 1552-1563. DOI: 10.1002/1529-0131(199809)41:9<1552::AID-ART5>3.0.CO;2-W

6. Weinblatt, M.E., Kremer, J.M., Bankhurst, A.D., Bulpitt, K.J., Fleischmann, R.M., Fox, R.I., et al. (1999) A trial of etanercept, a recombinant tumor necrosis factor receptor:Fc fusion protein, in patients with rheumatoid arthritis receiving methotrexate. New England Journal of Medicine, 340(4): 253-259. DOI: 10.1056/NEJM199901283400401

7. Keystone, E.C., Kavanaugh, A.F., Sharp, J.T., Tannenbaum, H., Hua, Y., Teoh, L.S., et al. (2004) Radiographic, clinical, and functional outcomes of treatment with adalimumab plus methotrexate. Arthritis & Rheumatism, 50(5): 1400-1411. DOI: 10.1002/art.20217

8. Smolen, J.S., Landewe, R., Breedveld, F.C., Dougados, M., Emery, P., Gaujoux-Viala, C., et al. (2010) EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs. Annals of the Rheumatic Diseases, 69(6): 964-975. DOI: 10.1136/ard.2009.126532

9. Singh, J.A., Saag, K.G., Bridges, S.L., Akl, E.A., Bannuru, R.R., Sullivan, M.C., et al. (2016) 2015 American College of Rheumatology guideline for the treatment of rheumatoid arthritis. Arthritis Care & Research, 68(1): 1-25. DOI: 10.1002/acr.22783

10. Smolen, J.S., Landewe, R.B.M., Bijlsma, J.W.J., Burmester, G.R., Dougados, M., Kerschbaumer, A., et al. (2020) EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2019 update. Annals of the Rheumatic Diseases, 79(6): 685-699. DOI: 10.1136/annrheumdis-2019-216655

11. Fraenkel, L., Bathon, J.M., England, B.R., St Clair, E.W., Arayssi, T., Carandang, K., et al. (2021) 2021 American College of Rheumatology guideline for the treatment of rheumatoid arthritis. Arthritis & Rheumatology, 73(7): 1108-1123. DOI: 10.1002/art.41752

12. Genovese, M.C., Bathon, J.M., Fleischmann, R.M., Moreland, L.W., Martin, R.W., Whitmore, J.B., et al. (2005) Longterm safety, efficacy, and radiographic outcome with etanercept treatment in patients with early rheumatoid arthritis. Journal of Rheumatology, 32(7): 1232-1242.

13. Klareskog, L., van der Heijde, D., de Jager, J.P., Gough, A., Kalden, J., Malaise, M., et al. (2004) Therapeutic effect of the combination of etanercept and methotrexate compared with each treatment alone in patients with rheumatoid arthritis. Lancet, 363(9410): 675-681. DOI: 10.1016/S0140-6736(04)15640-7

14. Breedveld, F.C., Weisman, M.H., Kavanaugh, A.F., Cohen, S.B., Pavelka, K., van Vollenhoven, R., et al. (2006) The PREMIER study: a multicenter, randomized, double-blind clinical trial comparing combination therapy with adalimumab plus methotrexate versus methotrexate alone or adalimumab alone in patients with early, aggressive rheumatoid arthritis. Arthritis & Rheumatism, 54(1): 26-37. DOI: 10.1002/art.21519

15. Emery, P., Fleischmann, R., Filipowicz-Sosnowska, A., Schechtman, J., Szczepanski, L., Kavanaugh, A., et al. (2008) The efficacy and safety of rituximab in patients with active rheumatoid arthritis despite methotrexate treatment: results of a phase IIb randomized, double-blind, placebo-controlled, dose-ranging trial. Arthritis & Rheumatism, 54(5): 1390-1400. DOI: 10.1002/art.21778

16. Cohen, S.B., Emery, P., Greenwald, M.W., Dougados, M., Furie, R.A., Genovese, M.C., et al. (2006) Rituximab for rheumatoid

© 2022 Somato Publications. All rights reserved.

arthritis refractory to anti-tumor necrosis factor therapy. Arthritis & Rheumatism, 54(9): 2793-2806. DOI: 10.1002/art.22025

17. Kremer, J.M., Genant, H.K., Moreland, L.W., Russell, A.S., Emery, P., Abud-Mendoza, C., et al. (2008) Effects of abatacept in patients with methotrexate-resistant active rheumatoid arthritis. Annals of Internal Medicine, 144(12): 865-876. DOI: 10.7326/0003-4819-144-12-200606200-00003

18. Genovese, M.C., Becker, J.C., Schiff, M., Luggen, M., Sherrer, Y., Kremer, J., et al. (2005) Abatacept for rheumatoid arthritis refractory to tumor necrosis factor alpha inhibition. New England Journal of Medicine, 353(11): 1114-1123. DOI: 10.1056/NEJMoa050524

19. Jones, G., Sebba, A., Gu, J., Lowenstein, M.B., Calvo, A., Gomez-Reino, J.J., et al. (2010) Comparison of tocilizumab monotherapy versus methotrexate monotherapy in patients with moderate to severe rheumatoid arthritis: the AMBITION study. Annals of the Rheumatic Diseases, 69(1): 88-96. DOI: 10.1136/ard.2008.105197

20. Emery, P., Keystone, E., Tony, H.P., Cantagrel, A., van Vollenhoven, R., Sanchez, A., et al. (2008) IL-6 receptor inhibition with tocilizumab improves treatment outcomes in patients with rheumatoid arthritis refractory to anti-tumour necrosis factor biologicals. Annals of the Rheumatic Diseases, 67(11): 1516-1523. DOI: 10.1136/ard.2008.092932

21. Gottenberg, J.E., Brocq, O., Perdriger, A., Lassoued, S., Berthelot, J.M., Wendling, D., et al. (2016) Non-TNF-targeted biologic vs a second anti-TNF drug to treat rheumatoid arthritis in patients with insufficient response to a first anti-TNF drug. JAMA, 316(11): 1172-1180. DOI: 10.1001/jama.2016.13512

22. Keystone, E., Heijde, D., Mason, D., Landewe, R., Vollenhoven, R.V., Combe, B., et al. (2008) Certolizumab pegol plus methotrexate is significantly more effective than placebo plus methotrexate in active rheumatoid arthritis: findings of a fifty-two-week, phase III, multicenter, randomized, double-blind, placebo-controlled, parallel-group study. Arthritis & Rheumatism, 58(11): 3319-3329. DOI: 10.1002/art.23964

23. Markusse, I.M., Akdemir, G., Dirven, L., Goekoop-Ruiterman, Y.P.M., van Groenendael, J.H.L.M., Han, K.H., et al. (2016) Long-term outcomes of patients with recent-onset rheumatoid arthritis after 10 years of tight controlled treatment: a randomized trial. Annals of Internal Medicine, 164(8): 523-531. DOI: 10.7326/M15-0919

24. Radner, H., Smolen, J.S., Aletaha, D. (2015) Remission in rheumatoid arthritis: benefit over low disease activity in patient-reported outcomes and costs. Arthritis Research & Therapy, 17: 251. DOI: 10.1186/s13075-015-0764-3

25. Felson, D.T., Smolen, J.S., Wells, G., Zhang, B., van Tuyl, L.H.D., Funovits, J., et al. (2011) American College of Rheumatology/European League Against Rheumatism provisional definition of remission in rheumatoid arthritis for clinical trials. Arthritis & Rheumatism, 63(3): 573-586. DOI: 10.1002/art.30129

26. Finckh, A., Simard, J.F., Gabay, C., Guerne, P.A. (2006) Evidence for differential acquired drug resistance to anti-tumour necrosis factor agents in rheumatoid arthritis. Annals of the Rheumatic Diseases, 65(6): 746-752. DOI: 10.1136/ard.2005.045062

27. Hyrich, K.L., Watson, K.D., Silman, A.J., Symmons, D.P.M. (2006) Predictors of response to anti-TNF-alpha therapy among patients with rheumatoid arthritis: results from the British Society for Rheumatology Biologics Register. Rheumatology, 45(12): 1558-1565. DOI: 10.1093/rheumatology/kel149

28. Hetland, M.L., Christensen, I.J., Tarp, U., Dreyer, L., Hansen, A., Hansen, I.T., et al. (2010) Direct comparison of treatment responses, remission rates, and drug adherence in patients with rheumatoid arthritis treated with adalimumab, etanercept, or infliximab. Arthritis & Rheumatism, 62(1):

© 2022 Somato Publications. All rights reserved.

22-32. DOI: 10.1002/art.27227

29. Bongartz, T., Sutton, A.J., Sweeting, M.J., Buchan, I., Matteson, E.L., Montori, V. (2006) Anti-TNF antibody therapy in rheumatoid arthritis and the risk of serious infections and malignancies. JAMA, 295(19): 2275-2285. DOI: 10.1001/jama.295.19.2275

30. Listing, J., Strangfeld, A., Kary, S., Rau, R., von Hinueber, U., Stoyanova-Scholz, M., et al. (2005) Infections in patients with rheumatoid arthritis treated with biologic agents. Arthritis & Rheumatism, 52(11): 3403-3412. DOI: 10.1002/art.21386

31. Galloway, J.B., Hyrich, K.L., Mercer, L.K., Dixon, W.G., Fu, B., Ustianowski, A.P., et al. (2011) Anti-TNF therapy is associated with an increased risk of serious infections in patients with rheumatoid arthritis especially in the first 6 months of treatment. Rheumatology, 50(1): 124-131. DOI: 10.1093/rheumatology/keq242

32. Singh, J.A., Wells, G.A., Christensen, R., Tanjong Ghogomu, E., Maxwell, L., Macdonald, J.K., et al. (2011) Adverse effects of biologics: a network meta-analysis and Cochrane overview. Cochrane Database of Systematic Reviews, (2): CD008794. DOI: 10.1002/14651858.CD008794.pub2

© 2022 Somato Publications. All rights reserved.