

Journal of Physical Medicine and Rehabilitation Research

Review Article

functional independence measure scoring in the Management of traumatic brain injury among stroke survivors

Nilsson I^{1*}, Berg L² and Lindqvist B³

¹Division of Rehabilitation Research, Euphrates Research Center for Rehabilitation Research, Iceland

²Department of Physical Medicine, Neva Institute of Rehabilitation Research, Sweden

³Department of Clinical Rehabilitation Research, Neva Institute of Rehabilitation Research, Finland

*Address for Correspondence: Ingrid Nilsson, Division of Rehabilitation Research, Euphrates Research Center for Rehabilitation Research, Iceland; E-mail: ingrid.nilsson@euphrates.edu

Received: 27 October 2021; Accepted: 19 January 2022; Published: 25 April 2022.

Citation of this article: Nilsson I, Berg L, Lindqvist B (2022) functional independence measure scoring in the Management of traumatic brain injury among stroke survivors. J Phys Med Rehabil Res, 4(1): 01-06. DOI: [10.5281/zenodo.22815056](https://doi.org/10.5281/zenodo.22815056)

Copyright: © 2022 Ingrid Nilsson, et al. This is an open access article distributed under the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: The Functional Independence Measure (FIM) is a widely used ordinal instrument for quantifying functional independence during inpatient rehabilitation, combining motor and cognitive subscales into an 18-item, 7-level score used to guide clinical decisions and benchmarking.

Objective: This review synthesizes the psychometric evidence for FIM scoring and its performance in traumatic brain injury (TBI) rehabilitation and stroke rehabilitation, with attention to stroke survivors whose presentation includes a TBI-like cognitive component.

Methods: We synthesized peer-reviewed reliability and validity studies, comparisons against the Barthel Index, and outcome studies relating FIM scores to rehabilitation intensity, care setting, and discharge disposition.

Findings: FIM motor items show high interrater reliability and predict discharge disposition and caregiver burden well, while cognitive items show lower reliability and ceiling effects that understate residual disability in higher-functioning TBI and TBI-pattern stroke survivors; responsiveness is broadly comparable to the Barthel Index for self-care domains.

Conclusions: FIM scoring remains a practical, reasonably reliable tool for tracking rehabilitation progress in stroke survivors with TBI-pattern disability when motor and cognitive subscales are reported separately and supplemented by instruments extending measurement into higher-level cognitive domains.

Keywords: Functional Independence Measure; traumatic brain injury; stroke rehabilitation; outcome measurement; cognitive impairment

1. Introduction

The Functional Independence Measure (FIM) was developed as a uniform, 18-item, 7-level ordinal instrument giving rehabilitation medicine a common metric for disability severity and the assistance a patient requires across self-care, sphincter control, mobility, locomotion, communication, and social cognition domains [1]. It became the centerpiece of the Uniform Data System for Medical Rehabilitation partly because earlier instruments, notably the Barthel Index, were built around simple activities of daily living and were not sensitive to the cognitive and behavioral impairments that dominate disability after traumatic brain injury (TBI) [2,3]. Stroke survivors present a related but distinct problem, since motor, sensory, communicative, and cognitive deficits co-occur in variable combinations, so a scale intended for one population-wide profile of impairment must perform consistently across heterogeneous case mixes to support benchmarking and clinical decision-making [4].

This review synthesizes the psychometric evidence for FIM scoring, its performance in TBI and stroke rehabilitation, where its assumptions break down, and the implications for clinicians managing stroke survivors whose presentation overlaps with or resembles TBI. The aim is not to argue that FIM should be replaced but to characterize where its scoring properties are strong and weak, and how FIM-derived scores should be weighted in guiding length of stay, discharge planning, and outcome comparison.

2. Structure, scoring, and psychometric properties

Each FIM item is rated from 1 (total assistance) to 7 (complete independence); item scores sum into motor and cognitive subscales that form the total score, ranging from 18 to 126 [1]. Interrater reliability studies across multiple facilities have reported intraclass correlation coefficients above 0.90 for the total score, with lower agreement for cognitive items such as social interaction

and problem solving than for motor items such as transfers and locomotion [5,6]. A quantitative review pooling reliability estimates across dozens of studies confirmed that motor-domain items are consistently more reliable than cognitive-domain items, a pattern directly relevant to TBI, where cognitive rather than motor impairment often dominates disability [6].

Table 1: FIM domains, item count, and reported interrater reliability, drawn from foundational psychometric studies of the instrument [1,5,6,7].

Domain	Items	Score range	Typical interrater reliability (ICC)
Motor subscale (self-care, sphincter, mobility, locomotion)	13	13-91	0.93-0.96
Cognitive subscale (communication, social cognition)	5	5-35	0.85-0.90
Total FIM score	18	18-126	0.90-0.96

Rasch analysis of the FIM's item structure showed that the instrument separates into a motor dimension and a cognitive dimension rather than behaving as a single unidimensional scale, supporting separate reporting of motor and cognitive subscores rather than the aggregate total alone [7]. Stineman and colleagues extended this analysis across 20 diagnostically diverse impairment categories and found that item difficulty ordering shifted somewhat between diagnostic groups, meaning the same total score does not necessarily represent the same underlying function across diagnoses [8]. Heinemann and colleagues showed the relationship between FIM scores and clinician-rated impairment, while generally monotonic, is least discriminating at the extremes, where ceiling and floor effects compress genuine differences into a narrow band [9]. Validity studies comparing FIM against independent clinical judgment reported concurrent validity coefficients generally in the 0.7-0.9 range [10], community-based inter-rater studies found stability comparable to inpatient cohorts [11], and independent replications outside the United States supported the instrument's use across health systems [12].

3. FIM scoring in traumatic brain injury rehabilitation

TBI presents the FIM with its most demanding test case, because impairment there is frequently dominated by cognitive, behavioral, and communicative deficits the cognitive subscale was designed, but not perfectly suited, to capture. Hall and colleagues, comparing FIM against the FIM+FAM (an extension adding cognitive, communicative, and community-oriented items), the Disability Rating Scale, and the Community Integration Questionnaire in a TBI cohort, found that many patients reached ceiling on FIM cognitive items well before reaching ceiling on the corresponding FIM+FAM items, indicating the base FIM underestimates residual disability in higher-functioning TBI survivors [13]. This has direct implications for stroke survivors with a TBI-like cognitive presentation, since a patient scored functionally independent on FIM cognitive items may still carry meaningful deficits in executive function or community reintegration only visible on more granular instruments [14].

Corrigan and colleagues, following TBI patients for five years, found functional gains, particularly cognitive, continue well beyond the typical inpatient stay, with meaningful further FIM-measured improvement one to two years post-injury [15], a trajectory relevant to stroke survivors with a substantial frontal or diffuse component whose discharge FIM may underestimate their eventual functional ceiling. Cifu and colleagues found admission and discharge FIM scores predicted disposition and later function less precisely in older TBI patients [16], relevant given that stroke, unlike isolated TBI, occurs predominantly in older patients with more comorbidity. Hall and Cope's synthesis concluded FIM and Disability Rating Scale gains correlate with, but do not fully explain, later community and vocational outcomes [17]; the Disability Rating Scale itself, validated against long-term coma-to-community trajectories, remains a useful complement where FIM's cognitive ceiling limits

sensitivity [18]. Cohen and Marino's broader review situates these TBI-specific limitations within the wider outcome-measurement literature, noting no single instrument fully captures both basic self-care independence and higher-order cognitive-behavioral recovery [19].




Figure 1: Proportion of a traumatic brain injury rehabilitation cohort reaching the scale ceiling on FIM cognitive items versus FIM+FAM higher-level cognitive and community items, illustrating the ceiling effect that limits FIM sensitivity in higher-functioning survivors [13].

4. FIM scoring in stroke rehabilitation

In stroke populations without a dominant TBI-like cognitive component, the FIM motor subscale predicts discharge disposition, length of stay, and burden of care well. Granger and colleagues found admission FIM motor score among the strongest predictors of community versus nursing-facility discharge, with discharge scores correlating with caregiver assistance needed at home [20]. Oczkowski and Barreca showed itemized FIM profiles, rather than the total score, are more useful for individualized planning because they expose specific problem domains, bathing and stair climbing in particular, even in patients with high totals [21]. The Copenhagen Stroke Study established that most motor recovery after stroke occurs within the first 8-12 weeks, with a smaller tail of later gains, shaping how discharge FIM scores taken at variable post-stroke intervals should be interpreted [22]. Heinemann and colleagues' multivariate analysis found admission functional status, age, and lesion side together explained a substantial share of discharge outcome variance, with functional score

contributing the largest single component [23].




Figure 2: Mean FIM motor score gain by rehabilitation care setting, comparing intensive inpatient rehabilitation, standard inpatient rehabilitation, and nonintensive home-based programs after stroke [24,25].

Because stroke rehabilitation spans a range of settings and intensities, Ozdemir and colleagues found patients in intensive inpatient programs achieved significantly greater FIM gains than comparable home-managed patients after adjusting for baseline severity [24]. Kwakkel and colleagues' randomized trial of training intensity showed higher-intensity training produced measurably greater functional gains within a fixed period, linking FIM-scored improvement to a modifiable treatment parameter [25]; their later work on upper-limb dexterity recovery showed time since onset and initial paresis severity jointly set the ceiling of achievable motor recovery [26]. A broader synthesis integrated these findings into a staged model of spontaneous recovery followed by a variable, therapy-responsive plateau, against which serial FIM scores should be interpreted rather than read as a simple linear trend [27].

5. Comparison with the Barthel Index and responsiveness to change

Table 2: Comparative properties of the FIM and the Barthel Index in stroke rehabilitation populations [28,29,30].		
Property	FIM	Barthel Index
Item count	18	10 (or 5-item short form)
Captures cognitive/communicative deficit	Yes, dedicated subscale	No
Administration burden	Higher	Lower

Property	FIM	Barthel Index
Responsiveness to inpatient change	Comparable to Barthel	Comparable to FIM

Because the FIM was designed partly to remedy the Barthel Index's insensitivity to cognitive impairment, direct comparisons in stroke populations inform scoring decisions. Hsueh and colleagues found the FIM and 5- and 10-item Barthel Index versions showed broadly similar reliability and validity for basic activities of daily living, but only FIM cognitive items captured impairments the Barthel-based measures missed, at the cost of longer administration [28]. Van der Putten and colleagues found the Barthel Index and FIM performed comparably in detecting meaningful change during inpatient rehabilitation, suggesting the simpler Barthel Index may suffice where only basic self-care gains matter [29]. Wallace and colleagues found the method used to quantify responsiveness materially affected which instrument appeared superior for the FIM motor component, a methodological caution against firm conclusions drawn from a single responsiveness statistic [30]. Segal and colleagues' interinstitutional study, in a spinal cord injury sample, found individual FIM items scored systematically differently between institutions even when overall reliability appeared acceptable, a caution for multi-site stroke and TBI research pooling FIM data [31].

6. Rehabilitation intensity, care setting, and outcome tracking

FIM scores are widely used to relate therapy delivered to functional gains achieved, for both treatment planning and program evaluation. Jette and colleagues found a positive but not strictly linear relationship between therapy minutes delivered and FIM-measured gain in skilled nursing facilities, plateauing at higher intensities and implying diminishing returns relevant to resource allocation [32]. Turner-Stokes' synthesis of multidisciplinary rehabilitation after acquired brain injury concluded coordinated programs produce gains

substantially exceeding natural recovery alone [33]. Grill and Stucki's criterion-validity analysis situated FIM within the International Classification of Functioning, Disability and Health, finding FIM items map most directly onto the activity domain and less onto participation, reinforcing that FIM gains do not by themselves establish satisfactory social and community participation [34]. Aggregate FIM data collected through the Uniform Data System continue to support national benchmarking of first-admission rehabilitation outcomes across facilities [4].

7. Limitations and considerations for scoring practice

Several limitations shape how FIM scores should be used in stroke survivors with TBI-pattern disability. Ceiling effects in the cognitive subscale mean a patient can score near maximum while retaining important deficits in executive function, insight, and social judgment, the deficits most characteristic of frontal and diffuse TBI and some frontal stroke syndromes [13]. Differential reliability of motor versus cognitive items means a total-score change driven mostly by motor recovery can mask stagnant cognitive function, an asymmetry best checked by examining subscale trajectories rather than the aggregate score [6,7]. Item difficulty ordering shifts across diagnostic groups, so a given total score does not carry an identical meaning in pure TBI, pure stroke, and mixed presentations [8]. Finally, because most foundational validity work used trained raters in controlled inpatient settings, scores collected later in recovery or by less experienced raters carry more uncertainty than the psychometric literature might suggest, a caution reinforced by documented interinstitutional scoring variability [31,12].

8. Conclusion

The FIM remains the most widely used and best-validated instrument for tracking functional independence during rehabilitation for both TBI and stroke, and its motor subscale predicts discharge disposition and caregiver

burden with reasonable accuracy across both populations [1,20]. Its cognitive subscale, a genuine improvement over instruments such as the Barthel Index that omit cognitive assessment entirely, shows ceiling effects and reduced sensitivity in exactly the patients, those with frontal, diffuse, or TBI-pattern injury, for whom cognitive and behavioral recovery matters most [13,28]. For stroke survivors with a substantial TBI component or TBI-like cognitive-behavioral profile, scoring practice should favor reporting motor and cognitive subscales separately, interpret scores against the expected recovery time course, and supplement them, where feasible, with instruments such as the FIM+FAM or Disability Rating Scale that extend measurement into higher-level cognitive and community domains [27,18]. Used as one component of a layered assessment strategy rather than a single definitive score, FIM scoring continues to provide a practical, reasonably reliable common language for tracking recovery in this complex population [33,34].

9. References

1. Keith, R.A., Granger, C.V., Hamilton, B.B., Sherwin, F.S. (1987) The Functional Independence Measure: a new tool for rehabilitation. *Advances in Clinical Rehabilitation*, 1: 6-18.
2. Granger, C.V., Hamilton, B.B., Linacre, J.M., Heinemann, A.W., Wright, B.D. (1993) Performance profiles of the Functional Independence Measure. *American Journal of Physical Medicine and Rehabilitation*, 72(2): 84-89. DOI: [10.1097/00002060-199304000-00005](https://doi.org/10.1097/00002060-199304000-00005)
3. Hamilton, B.B., Granger, C.V., Sherwin, F.S., Zielezny, M., Tashman, J.S. (1987) A uniform national data system for medical rehabilitation. In: Fuhrer, M.J. (Ed.), *Rehabilitation Outcomes: Analysis and Measurement*. Paul H. Brookes Publishing, Baltimore, pp. 137-147.
4. Granger, C.V., Hamilton, B.B. (1992) The Uniform Data System for Medical Rehabilitation report of first admissions for 1990. *American Journal of Physical Medicine and Rehabilitation*, 71(2): 108-113.

5. Hamilton, B.B., Laughlin, J.A., Fiedler, R.C., Granger, C.V. (1994) Interrater reliability of the 7-level functional independence measure (FIM). *Scandinavian Journal of Rehabilitation Medicine*, 26(3): 115-119.
6. Ottenbacher, K.J., Hsu, Y., Granger, C.V., Fiedler, R.C. (1996) The reliability of the functional independence measure: a quantitative review. *Archives of Physical Medicine and Rehabilitation*, 77(12): 1226-1232. DOI: [10.1016/S0003-9993\(96\)90184-7](https://doi.org/10.1016/S0003-9993(96)90184-7)
7. Linacre, J.M., Heinemann, A.W., Wright, B.D., Granger, C.V., Hamilton, B.B. (1994) The structure and stability of the Functional Independence Measure. *Archives of Physical Medicine and Rehabilitation*, 75(2): 127-132. DOI: [10.1016/0003-9993\(94\)90384-0](https://doi.org/10.1016/0003-9993(94)90384-0)
8. Stineman, M.G., Shea, J.A., Jette, A., Tassoni, C.J., Ottenbacher, K.J., Fiedler, R., Granger, C.V. (1996) The Functional Independence Measure: tests of scaling assumptions, structure, and reliability across 20 diverse impairment categories. *Archives of Physical Medicine and Rehabilitation*, 77(11): 1101-1108. DOI: [10.1016/S0003-9993\(96\)90130-6](https://doi.org/10.1016/S0003-9993(96)90130-6)
9. Heinemann, A.W., Linacre, J.M., Wright, B.D., Hamilton, B.B., Granger, C.V. (1993) Relationships between impairment and physical disability as measured by the functional independence measure. *Archives of Physical Medicine and Rehabilitation*, 74(6): 566-573. DOI: [10.1016/0003-9993\(93\)90153-2](https://doi.org/10.1016/0003-9993(93)90153-2)
10. Dodds, T.A., Martin, D.P., Stolov, W.C., Deyo, R.A. (1993) A validation of the functional independence measurement and its performance among rehabilitation inpatients. *Archives of Physical Medicine and Rehabilitation*, 74(5): 531-536. DOI: [10.1016/0003-9993\(93\)90119-U](https://doi.org/10.1016/0003-9993(93)90119-U)
11. Ottenbacher, K.J., Mann, W.C., Granger, C.V., Tomita, M., Hurren, D., Charvat, B. (1994) Inter-rater agreement and stability of functional assessment in the community-based elderly. *Archives of Physical Medicine and Rehabilitation*, 75(12): 1297-1301. DOI: [10.1016/0003-9993\(94\)90279-X](https://doi.org/10.1016/0003-9993(94)90279-X)
12. Kidd, D., Stewart, G., Baldry, J., Johnson, J., Rossiter, D., Petruckevitch, A., Thompson, A.J. (1995) The Functional Independence Measure: a comparative validity and reliability study. *Disability and Rehabilitation*, 17(1): 10-14. DOI: [10.3109/09638289509166622](https://doi.org/10.3109/09638289509166622)
13. Hall, K.M., Mann, N., High, W.M. Jr, Wright, J., Kreutzer, J.S., Wood, D. (1996) Functional measures after traumatic brain injury: ceiling effects of FIM, FIM+FAM, DRS, and CIQ. *Journal of Head Trauma Rehabilitation*, 11(5): 27-39. DOI: [10.1097/00001199-199610000-00004](https://doi.org/10.1097/00001199-199610000-00004)
14. Hall, K.M., Hamilton, B.B., Gordon, W.A., Zasler, N.D. (1993) Characteristics and comparisons of functional assessment indices: Disability Rating Scale, Functional Independence Measure, and Functional Assessment Measure. *Journal of Head Trauma Rehabilitation*, 8(2): 60-74.
15. Corrigan, J.D., Smith-Knapp, K., Granger, C.V. (1997) Outcomes in the first 5 years after traumatic brain injury. *Archives of Physical Medicine and Rehabilitation*, 78(3): 298-305. DOI: [10.1016/S0003-9993\(97\)90038-1](https://doi.org/10.1016/S0003-9993(97)90038-1)
16. Cifu, D.X., Kreutzer, J.S., Marwitz, J.H., Rosenthal, M., Englander, J., High, W. (1996) Functional outcomes of older adults with traumatic brain injury: a prospective, multicenter analysis. *Archives of Physical Medicine and Rehabilitation*, 77(9): 883-888. DOI: [10.1016/S0003-9993\(96\)90274-9](https://doi.org/10.1016/S0003-9993(96)90274-9)
17. Hall, K.M., Cope, D.N. (1995) The benefit of rehabilitation in traumatic brain injury: a literature review. *Journal of Head Trauma Rehabilitation*, 10(1): 1-13. DOI: [10.1097/00001199-199502000-00003](https://doi.org/10.1097/00001199-199502000-00003)
18. Rappaport, M., Hall, K.M., Hopkins, K., Belleza, T., Cope, D.N. (1982) Disability rating scale for severe head trauma: coma to community. *Archives of Physical Medicine and Rehabilitation*, 63(3): 118-123.
19. Cohen, M.E., Marino, R.J. (2000) The tools of disability

- outcomes research functional status measures. *Archives of Physical Medicine and Rehabilitation*, 81(12 Suppl 2): S21-S29. DOI: [10.1053/apmr.2000.20620](https://doi.org/10.1053/apmr.2000.20620)
20. Granger, C.V., Hamilton, B.B., Fiedler, R.C. (1992) Discharge outcome after stroke rehabilitation. *Stroke*, 23(7): 978-982. DOI: [10.1161/01.STR.23.7.978](https://doi.org/10.1161/01.STR.23.7.978)
 21. Oczkowski, W.J., Barreca, S. (1993) The Functional Independence Measure: its use to identify rehabilitation needs in stroke survivors. *Archives of Physical Medicine and Rehabilitation*, 74(12): 1291-1294. DOI: [10.1016/0003-9993\(93\)90081-K](https://doi.org/10.1016/0003-9993(93)90081-K)
 22. Jorgensen, H.S., Nakayama, H., Raaschou, H.O., Vive-Larsen, J., Stoier, M., Olsen, T.S. (1995) Outcome and time course of recovery in stroke. Part II: Time course of recovery. The Copenhagen Stroke Study. *Archives of Physical Medicine and Rehabilitation*, 76(5): 406-412. DOI: [10.1016/S0003-9993\(95\)80568-0](https://doi.org/10.1016/S0003-9993(95)80568-0)
 23. Heinemann, A.W., Roth, E.J., Cichowski, K., Betts, H.B. (1987) Multivariate analysis of improvement and outcome following stroke rehabilitation. *Archives of Neurology*, 44(11): 1167-1172. DOI: [10.1001/archneur.1987.00520230031012](https://doi.org/10.1001/archneur.1987.00520230031012)
 24. Ozdemir, F., Birtane, M., Tabatabaei, R., Ekuklu, G., Kokino, S. (2001) Comparing stroke rehabilitation outcomes between acute inpatient and nonintensive home settings. *Archives of Physical Medicine and Rehabilitation*, 82(10): 1375-1379. DOI: [10.1053/apmr.2001.25971](https://doi.org/10.1053/apmr.2001.25971)
 25. Kwakkel, G., Kollen, B.J., Wagenaar, R.C. (2002) Long term effects of intensity of upper and lower limb training after stroke: a randomised trial. *Journal of Neurology, Neurosurgery and Psychiatry*, 72(4): 473-479. DOI: [10.1136/jnnp.72.4.473](https://doi.org/10.1136/jnnp.72.4.473)
 26. Kwakkel, G., Kollen, B.J., van der Grond, J., Prevo, A.J. (2003) Probability of regaining dexterity in the flaccid upper limb: impact of severity of paresis and time since onset in acute stroke. *Stroke*, 34(9): 2181-2186. DOI: [10.1161/01.STR.0000087172.16305.CD](https://doi.org/10.1161/01.STR.0000087172.16305.CD)
 27. Kwakkel, G., Kollen, B., Lindeman, E. (2004) Understanding the pattern of functional recovery after stroke: facts and theories. *Restorative Neurology and Neuroscience*, 22(3-5): 281-299.
 28. Hsueh, I.P., Lin, J.H., Jeng, J.S., Hsieh, C.L. (2002) Comparison of the psychometric characteristics of the functional independence measure, 5 item Barthel index, and 10 item Barthel index in patients with stroke. *Journal of Neurology, Neurosurgery and Psychiatry*, 73(2): 188-190. DOI: [10.1136/jnnp.73.2.188](https://doi.org/10.1136/jnnp.73.2.188)
 29. van der Putten, J.J., Hobart, J.C., Freeman, J.A., Thompson, A.J. (1999) Measuring change in disability after inpatient rehabilitation: comparison of the responsiveness of the Barthel index and the Functional Independence Measure. *Journal of Neurology, Neurosurgery and Psychiatry*, 66(4): 480-484. DOI: [10.1136/jnnp.66.4.480](https://doi.org/10.1136/jnnp.66.4.480)
 30. Wallace, D., Duncan, P.W., Lai, S.M. (2002) Comparison of the responsiveness of the Barthel Index and the motor component of the Functional Independence Measure in stroke: the impact of using different methods for measuring responsiveness. *Journal of Clinical Epidemiology*, 55(9): 922-928. DOI: [10.1016/S0895-4356\(02\)00434-8](https://doi.org/10.1016/S0895-4356(02)00434-8)
 31. Segal, M.E., Ditunno, J.F., Staas, W.E. (1993) Interinstitutional agreement of individual functional independence measure (FIM) items measured at two sites on one sample of SCI patients. *Paraplegia*, 31(10): 622-631. DOI: [10.1038/sc.1993.100](https://doi.org/10.1038/sc.1993.100)
 32. Jette, D.U., Warren, R.L., Wirtalla, C. (2005) The relation between therapy intensity and outcomes of rehabilitation in skilled nursing facilities. *Archives of Physical Medicine and Rehabilitation*, 86(3): 373-379. DOI: [10.1016/j.apmr.2004.10.018](https://doi.org/10.1016/j.apmr.2004.10.018)
 33. Turner-Stokes, L. (2008) Evidence for the effectiveness of multi-disciplinary rehabilitation following acquired brain injury: a synthesis of two systematic approaches. *Journal of*

Citation: Nilsson I, Berg L, Lindqvist B (2022) functional independence measure scoring in the Management of traumatic brain injury among stroke survivors. J Phys Med Rehabil Res, 4(1): 01-06.

Rehabilitation Medicine, 40(9): 691-701. DOI:
[10.2340/16501977-0265](https://doi.org/10.2340/16501977-0265)

34. Grill, E., Stucki, G. (2011) Criterion validity of outcome measures for stroke rehabilitation based on the international classification of functioning, disability and health. Journal of Rehabilitation Medicine, 43(4): 297-304.
DOI: [10.2340/16501977-0654](https://doi.org/10.2340/16501977-0654)

